

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/003483

International filing date: 27 January 2005 (27.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/539,837  
Filing date: 28 January 2004 (28.01.2004)

Date of receipt at the International Bureau: 07 April 2005 (07.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1300910

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*April 01, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/539,837**

**FILING DATE: *January 28, 2004***

**RELATED PCT APPLICATION NUMBER: *PCT/US05/03483***



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office



012804

16805 U.S. PTO

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EV385630135US

INVENTOR(S)					
Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)			
Lori	Friedman	San Carlos, California			
Additional inventors are being named on the _____ 1 _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
MIGFRS AS MODIFIERS OF THE IGFR PATHWAY AND METHODS OF USE					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number: 23500					
OR					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		Zip	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages 80		<input type="checkbox"/> CD(s), Number _____			
<input type="checkbox"/> Drawing(s) Number of Sheets _____		<input checked="" type="checkbox"/> Other (specify) return postcard			
<input type="checkbox"/> Application Date Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.					FILING FEE Amount (\$)  160.00
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.					
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 50-1108					
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

[Page 1 of 2]

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME Laleh Shayesteh

TELEPHONE 650.837.8223

Date January 28, 2004

REGISTRATION NO. 47,937

(if appropriate)

Docket Number: EX04-007P

**USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

60/539837



012804

**Additional Page**

**Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.**

**Docket Number** EX04-007P

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any] )	Family or Surname	Residence (City and either State or Foreign Country)
Helen	Francis-Lang	San Francisco, California
Annette	Parks	Daly City, California
Kenneth	Shaw	
HaiGuang	Zhang	El Sobrante, California
Lynn Margaret	Bjerke	San Francisco, California
Monique	Nicoll	Pacific, California
Joanne I.	Adamkewicz	San Francisco, California

[Page 2 of 2]

Number 1 of 1

**WARNING:** Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

## MIGFRS AS MODIFIERS OF THE IGF1R PATHWAY AND METHODS OF USE

### BACKGROUND OF THE INVENTION

Somatic mutations in the PTEN (Phosphatase and Tensin homolog deleted on chromosome 10) gene are known to cause tumors in a variety of human tissues. In addition, germline mutations in PTEN are the cause of human diseases (Cowden disease and Bannayan-Zonana syndrome) associated with increased risk of breast and thyroid cancer (Nelen MR et al. (1997) Hum Mol Genet, 8:1383-1387; Liaw D et al. (1997) Nat Genet, 1:64-67; Marsh DJ et al. (1998) Hum Mol Genet, 3:507-515). PTEN is thought to act as a tumor suppressor by regulating several signaling pathways through the second messenger phosphatidylinositol 3,4,5 triphosphate (PIP3). PTEN dephosphorylates the D3 position of PIP3 and downregulates signaling events dependent on PIP3 levels (Maehama T and Dixon JE (1998) J Biol Chem, 273:13375-8). In particular, pro-survival pathways downstream of the insulin-like growth factor (IGF) pathway are regulated by PTEN activity. Stimulation of the IGF pathway, or loss of PTEN function, elevates PIP3 levels and activates pro-survival pathways associated with tumorigenesis (Stambolic V et al. (1998) Cell, 95:29-39). Consistent with this model, elevated levels of insulin-like growth factors I and II correlate with increased risk of cancer (Yu H et al (1999) J Natl Cancer Inst 91:151-156) and poor prognosis (Takanami I et al, 1996, J Surg Oncol 61(3):205-8). In addition, increased levels or activity of positive effectors of the IGF pathway, such as Akt and PI(3) kinase, have been implicated in several types of human cancer (Nicholson KM and Anderson NG (2002) Cellular Signalling, 14:381-395). In *Drosophila melanogaster*, as in vertebrates, the Insulin Growth Factor Receptor (IGFR) pathway includes the positive effectors PI(3) kinase, Akt, and PDK and the inhibitor, PTEN. These proteins have been implicated in multiple processes, including the regulation of cell growth and size as well as cell division and survival (Oldham S and Hafen E. (2003) Trends Cell Biol. 13:79-85; Garafolo RS. (2002) Trends Endocr. Metab. 13:156-162; Backman SA et al. (2002) Curr. Op. Neurobio. 12:1-7; Tapon N et al. (2001) Curr Op. Cell Biol. 13:731-737). Activation of the pathway in *Drosophila* can result in increases in cell size, cell number and organ size (Oldham S et al. (2002) Dev. 129:4103-

4109; Prober DA and Edgar BA. (2002) *Genes&Dev.* 16:2286-2299; Potter CJ et al. (2001) *Cell* 105:357-368; Verdu J et al. (1999) *Cell Biol.* 1:500-506).

The ability to manipulate the genomes of model organisms such as *Drosophila* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Mechler BM et al., 1985 *EMBO J* 4:1551-1557; Gateff E. 1982 *Adv. Cancer Res.* 37: 33-74; Watson KL., et al., 1994 *J Cell Sci.* 18: 19-33; Miklos GL, and Rubin GM. 1996 *Cell* 86:521-529; Wassarman DA, et al., 1995 *Curr Opin Gen Dev* 5: 44-50; and Booth DR. 1999 *Cancer Metastasis Rev.* 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a “genetic entry point”) that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a “modifier” involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as IGFR, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

## SUMMARY OF THE INVENTION

We have discovered genes that modify the IGFR pathway in *Drosophila*, and identified their human orthologs, hereinafter referred to as Modifier of IGFR (MIGFR). The invention provides methods for utilizing these IGFR modifier genes

and polypeptides to identify MIGFR-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired IGFR function and/or MIGFR function. Preferred MIGFR-modulating agents specifically bind to MIGFR polypeptides and restore IGFR function. Other preferred MIGFR-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress MIGFR gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

MIGFR modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with an MIGFR polypeptide or nucleic acid. In one embodiment, candidate MIGFR modulating agents are tested with an assay system comprising a MIGFR polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate IGFR modulating agents. The assay system may be cell-based or cell-free. MIGFR-modulating agents include MIGFR related proteins (e.g. dominant negative mutants, and biotherapeutics); MIGFR -specific antibodies; MIGFR -specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with MIGFR or compete with MIGFR binding partner (e.g. by binding to an MIGFR binding partner). In one specific embodiment, a small molecule modulator is identified using a binding assay. In specific embodiments, the screening assay system is selected from an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

In another embodiment, candidate IGFR pathway modulating agents are further tested using a second assay system that detects changes in the IGFR pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the IGFR pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the MIGFR function and/or the IGFR pathway in a mammalian cell by contacting the mammalian cell with

an agent that specifically binds a MIGFR polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated with the IGFR pathway.

### **DETAILED DESCRIPTION OF THE INVENTION**

A dominant loss of function screen was carried out in *Drosophila* to identify genes that interact with or modulate the IGFR signaling pathway. Modifiers of the IGFR pathway and their orthologs were identified. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, MIGFR genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective IGFR signaling pathway, such as cancer. Table 1 (Example II) lists the modifiers and their orthologs.

In vitro and in vivo methods of assessing MIGFR function are provided herein. Modulation of the MIGFR or their respective binding partners is useful for understanding the association of the IGFR pathway and its members in normal and disease conditions and for developing diagnostics and therapeutic modalities for IGFR related pathologies. MIGFR-modulating agents that act by inhibiting or enhancing MIGFR expression, directly or indirectly, for example, by affecting an MIGFR function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. MIGFR modulating agents are useful in diagnosis, therapy and pharmaceutical development.

#### **Nucleic acids and polypeptides of the invention**

Sequences related to MIGFR nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) or RefSeq number), shown in Table 1 and in the appended sequence listing.

The term “MIGFR polypeptide” refers to a full-length MIGFR protein or a functionally active fragment or derivative thereof. A “functionally active” MIGFR fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type MIGFR protein, such as antigenic or immunogenic activity,



enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of MIGFR proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active MIGFR polypeptide is a MIGFR derivative capable of rescuing defective endogenous MIGFR activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of an MIGFR, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). Methods for obtaining MIGFR polypeptides are also further described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of an MIGFR. In further preferred embodiments, the fragment comprises the entire functionally active domain.

The term "MIGFR nucleic acid" refers to a DNA or RNA molecule that encodes a MIGFR polypeptide. Preferably, the MIGFR polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human MIGFR. Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res

22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Drosophila*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term “orthologs” encompasses paralogs. As used herein, “percent (%) sequence identity” with respect to a subject sequence, or a specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. “Percent (%) amino acid sequence similarity” is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and

valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" ([www.psc.edu](http://www.psc.edu)) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of an MIGFR. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, *Current Protocol in Molecular Biology*, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of an MIGFR under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml

herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

### **Isolation, Production, Expression, and Mis-expression of MIGFR Nucleic Acids and Polypeptides**

MIGFR nucleic acids and polypeptides are useful for identifying and testing agents that modulate MIGFR function and for other applications related to the involvement of MIGFR in the IGFR pathway. MIGFR nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of

proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of an MIGFR protein for assays used to assess MIGFR function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) *Protein Expression: A Practical Approach*, Oxford University Press Inc., New York 1999; Stanbury PF et al., *Principles of Fermentation Technology*, 2<sup>nd</sup> edition, Elsevier Science, New York, 1995; Doonan S (ed.) *Protein Purification Protocols*, Humana Press, New Jersey, 1996; Coligan JE et al, *Current Protocols in Protein Science* (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant MIGFR is expressed in a cell line known to have defective IGFR function. The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding an MIGFR polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native MIGFR gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the MIGFR gene product, the expression vector can comprise a promoter operably linked to an MIGFR gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the MIGFR gene product based on the physical or functional properties of the MIGFR protein in *in vitro* assay systems (*e.g.* immunoassays).

The MIGFR protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the MIGFR gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native MIGFR proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of MIGFR or other genes associated with the IGFR pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

### **Genetically modified animals**

Animal models that have been genetically modified to alter MIGFR expression may be used in *in vivo* assays to test for activity of a candidate IGFR modulating agent, or to further assess the role of MIGFR in a IGFR pathway process such as apoptosis or cell proliferation. Preferably, the altered MIGFR expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal MIGFR expression. The genetically modified animal may additionally have altered IGFR expression (*e.g.* IGFR knockout). Preferred genetically modified animals are

mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, i.e. mosaic animals (see, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) or stably integrated into its germ line DNA (i.e., in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster et al., Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Pat. No. 4,873,191 by Wagner et al., and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sanford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a “knock-out” animal having a heterozygous or homozygous alteration in the sequence of an endogenous MIGFR gene that results in a decrease of MIGFR function, preferably such that MIGFR

expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse MIGFR gene is used to construct a homologous recombination vector suitable for altering an endogenous MIGFR gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, *Science* (1989) 244:1288-1292; Joyner *et al.*, *Nature* (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, *Science* (1989) 244:1281-1288; Simms *et al.*, *Bio/Technology* (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) *Scan J Immunol* 40:257-264; Declerck PJ *et al.*, (1995) *J Biol Chem.* 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the MIGFR gene, e.g., by introduction of additional copies of MIGFR, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the MIGFR gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, *PNAS* (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two



transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X et al (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the IGFR pathway, as animal models of disease and disorders implicating defective IGFR function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered MIGFR function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered MIGFR expression that receive candidate therapeutic agent.

In addition to the above-described genetically modified animals having altered MIGFR function, animal models having defective IGFR function (and otherwise normal MIGFR function), can be used in the methods of the present invention. For example, a IGFR knockout mouse can be used to assess, *in vivo*, the activity of a candidate IGFR modulating agent identified in one of the *in vitro* assays described below. Preferably, the candidate IGFR modulating agent when administered to a model system with cells defective in IGFR function, produces a detectable phenotypic change in the model system indicating that the IGFR function is restored, i.e., the cells exhibit normal cell cycle progression.

### **Modulating Agents**

The invention provides methods to identify agents that interact with and/or modulate the function of MIGFR and/or the IGFR pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the IGFR pathway, as well as in further analysis of the MIGFR protein and its contribution to the IGFR pathway. Accordingly,

the invention also provides methods for modulating the IGFR pathway comprising the step of specifically modulating MIGFR activity by administering a MIGFR-interacting or -modulating agent.

As used herein, an "MIGFR-modulating agent" is any agent that modulates MIGFR function, for example, an agent that interacts with MIGFR to inhibit or enhance MIGFR activity or otherwise affect normal MIGFR function. MIGFR function can be affected at any level, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the MIGFR - modulating agent specifically modulates the function of the MIGFR. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the MIGFR polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the MIGFR. These phrases also encompass modulating agents that alter the interaction of the MIGFR with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of an MIGFR, or to a protein/binding partner complex, and altering MIGFR function). In a further preferred embodiment, the MIGFR- modulating agent is a modulator of the IGFR pathway (e.g. it restores and/or upregulates IGFR function) and thus is also a IGFR-modulating agent.

Preferred MIGFR-modulating agents include small molecule compounds; MIGFR-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19<sup>th</sup> edition.

### **Small molecule modulators**

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight up to 10,000, preferably up to 5,000, more preferably up to

1,000, and most preferably up to 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the MIGFR protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for MIGFR-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the IGFR pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

### **Protein Modulators**

Specific MIGFR-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the IGFR pathway and related disorders, as well as in validation assays for other MIGFR-modulating agents. In a preferred embodiment, MIGFR-interacting proteins affect normal MIGFR function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, MIGFR-interacting proteins are useful in detecting and providing information about the function of MIGFR proteins, as is relevant to IGFR related disorders, such as cancer (e.g., for diagnostic means).

An MIGFR-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with an MIGFR, such as a member of the MIGFR pathway that modulates MIGFR expression, localization, and/or activity. MIGFR-modulators include dominant negative forms of MIGFR-interacting proteins and of MIGFR proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous MIGFR-interacting proteins (Finley, R. L. et al. (1996) in *DNA Cloning-Expression Systems: A Practical Approach*, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., *Gene* (2000) 250:1-14; Drees BL *Curr Opin Chem Biol* (1999) 3:64-70; Vidal M and Legrain P *Nucleic Acids Res* (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, *Nature* (2000) 405:837-846; Yates JR 3<sup>rd</sup>, *Trends Genet* (2000) 16:5-8).

An MIGFR-interacting protein may be an exogenous protein, such as an MIGFR-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory; Harlow and Lane (1999) *Using antibodies: a laboratory manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). MIGFR antibodies are further discussed below.

In preferred embodiments, an MIGFR-interacting protein specifically binds an MIGFR protein. In alternative preferred embodiments, an MIGFR-modulating agent binds an MIGFR substrate, binding partner, or cofactor.

### ***Antibodies***

In another embodiment, the protein modulator is an MIGFR specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify MIGFR modulators. The antibodies can also be used in dissecting the portions of the MIGFR pathway responsible for various cellular responses and in the general processing and maturation of the MIGFR.

Antibodies that specifically bind MIGFR polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of MIGFR polypeptide, and more preferably, to human MIGFR. Antibodies may be polyclonal,

monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of MIGFR which are particularly antigenic can be selected, for example, by routine screening of MIGFR polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Natl. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89; Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence of an MIGFR. Monoclonal antibodies with affinities of  $10^8 \text{ M}^{-1}$  preferably  $10^9 \text{ M}^{-1}$  to  $10^{10} \text{ M}^{-1}$ , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of MIGFR or substantially purified fragments thereof. If MIGFR fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of an MIGFR protein. In a particular embodiment, MIGFR-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of MIGFR-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding MIGFR polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to MIGFR polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al.,

Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

MIGFR-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be

produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

### ***Specific biotherapeutics***

In a preferred embodiment, an MIGFR-interacting protein may have biotherapeutic applications. Biotherapeutic agents formulated in pharmaceutically acceptable carriers and dosages may be used to activate or inhibit signal transduction pathways. This modulation may be accomplished by binding a ligand, thus inhibiting the activity of the pathway; or by binding a receptor, either to inhibit activation of, or to activate, the receptor. Alternatively, the biotherapeutic may itself be a ligand capable of activating or inhibiting a receptor. Biotherapeutic agents and methods of producing them are described in detail in U.S. Pat. No. 6,146,628.

When the MIGFR is a ligand, it may be used as a biotherapeutic agent to activate or inhibit its natural receptor. Alternatively, antibodies against MIGFR, as described in the previous section, may be used as biotherapeutic agents.

When the MIGFR is a receptor, its ligand(s), antibodies to the ligand(s) or the MIGFR itself may be used as biotherapeutics to modulate the activity of MIGFR in the IGFR pathway.

### **Nucleic Acid Modulators**

Other preferred MIGFR-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit MIGFR activity. Preferred nucleic acid modulators interfere with the function of the MIGFR nucleic acid such as DNA replication, transcription, translocation of the MIGFR RNA to the site of protein translation, translation of protein from the MIGFR RNA, splicing of the MIGFR RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the MIGFR RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to an MIGFR mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. MIGFR-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281;



Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred MIGFR nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al.*, Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, an MIGFR-specific nucleic acid modulator is used in an assay to further elucidate the role of the MIGFR in the IGFR pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, an MIGFR-specific antisense oligomer is used as a therapeutic agent for treatment of IGFR-related disease states.

### **Assay Systems**

The invention provides assay systems and screening methods for identifying specific modulators of MIGFR activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the MIGFR nucleic acid or protein. In general, secondary assays further assess the activity of a MIGFR modulating agent identified by a primary assay and may confirm that the modulating agent affects MIGFR in a manner relevant to the IGFR pathway. In some cases, MIGFR modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising an MIGFR polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. binding activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates MIGFR activity, and hence the IGFR pathway. The MIGFR polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

### **Primary Assays**

The type of modulator tested generally determines the type of primary assay.

#### ***Primary assays for small molecule modulators***

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or

crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding), transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of MIGFR and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when MIGFR-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the MIGFR protein may be assayed by various known methods such as substrate processing (*e.g.* ability of the candidate MIGFR-specific binding agents to function as negative effectors in MIGFR-expressing cells), binding equilibrium constants (usually at least about  $10^7 \text{ M}^{-1}$ , preferably at least about  $10^8 \text{ M}^{-1}$ , more preferably at least about  $10^9 \text{ M}^{-1}$ ), and immunogenicity (*e.g.* ability to elicit MIGFR specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a MIGFR polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The MIGFR polypeptide can be full length or a fragment thereof that retains functional MIGFR activity. The MIGFR polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The MIGFR polypeptide is preferably human MIGFR, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the

screening assay detects candidate agent-based modulation of MIGFR interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has MIGFR –specific binding activity, and can be used to assess normal MIGFR gene function.

Suitable assay formats that may be adapted to screen for MIGFR modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, *Curr Opin Chem Biol* (1998) 2:597-603; Sundberg SA, *Curr Opin Biotechnol* 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, *Nat Struct Biol* (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, *Curr Opin Chem Biol* (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate MIGFR and IGFR pathway modulators (*e.g.* U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); U.S. Pat. No. 6,114,132 (phosphatase and protease assays), U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

Protein kinases, key signal transduction proteins that may be either membrane-associated or intracellular, catalyze the transfer of gamma phosphate from adenosine triphosphate (ATP) to a serine, threonine or tyrosine residue in a protein substrate. Radioassays, which monitor the transfer from [ $\gamma$ - $^{32}\text{P}$  or  $^{33}\text{P}$ ]ATP, are frequently used to assay kinase activity. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from [ $\gamma$ - $^{33}\text{P}$ ] ATP to a biotinylated peptide substrate. The substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, *J Biomol Screen* (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from

free ligand. Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133). Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

Protein phosphatases catalyze the removal of a gamma phosphate from a serine, threonine or tyrosine residue in a protein substrate. Since phosphatases act in opposition to kinases, appropriate assays measure the same parameters as kinase assays. In one example, the dephosphorylation of a fluorescently labeled peptide substrate allows trypsin cleavage of the substrate, which in turn renders the cleaved substrate significantly more fluorescent (Nishikata M *et al.*, Biochem J (1999) 343:35-391). In another example, fluorescence polarization (FP), a solution-based, homogeneous technique requiring no immobilization or separation of reaction components, is used to develop high throughput screening (HTS) assays for protein phosphatases. This assay uses direct binding of the phosphatase with the target, and increasing concentrations of target-phosphatase increase the rate of dephosphorylation, leading to a change in polarization (Parker GJ et al., (2000) J Biomol Screen 5:77-88).

High-throughput assays, such as scintillation proximity assays, for synthase enzymes involved in fatty acid synthesis are known in the art (He X et al (2000) Anal Biochem 2000 Jun 15;282(1):107-14).

**Apoptosis assays.** Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of

fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the caspase cleavage activity as part of a cascade of events that occur during programmed cell death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-ONE<sup>TM</sup> Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death assay known to those skilled in the art, and available commercially (Roche, Cat# 1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses monoclonal antibodies directed against DNA and histones respectively, thus specifically determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due to the fact that DNA fragmentation occurs several hours before the plasma membrane breaks down, allowing for accumulation in the cytoplasm. Nucleosomes are not present in the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay system may comprise a cell that expresses an MIGFR, and that optionally has defective IGFR function (e.g. IGFR is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate IGFR modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate IGFR modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether MIGFR function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express MIGFR relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the MIGFR plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

**Cell proliferation and cell cycle assays.** Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specific to the phosphorylated form of the serine 10 residue of histone H3. (Chadlee, D.N. 1995, *J. Biol. Chem* 270:20098-105). Cell Proliferation may also be examined using [<sup>3</sup>H]-thymidine incorporation (Chen, J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [<sup>3</sup>H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL *et al.*, 1998, *In Vitro Cell Dev Biol Anim* 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt, MTS. MTS assays are commercially available, for example, the Promega CellTiter 96<sup>®</sup> Aqueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed with MIGFR are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Cell proliferation may also be assayed by measuring ATP levels as indicator of metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo™, which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) *Int J Radiat Biol Relat Stud Phys Chem Med* 49:237-55). Cells transfected with an MIGFR may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses an MIGFR, and that optionally has defective IGFR function (e.g. IGFR is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate IGFR modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate IGFR modulating agents that is initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test whether MIGFR function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express MIGFR relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the MIGFR plays a direct role in cell proliferation or cell cycle.

**Angiogenesis.** Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses an MIGFR, and that optionally has defective IGFR function (e.g. IGFR is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate IGFR modulating



agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate IGFR modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether MIGFR function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express MIGFR relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the MIGFR plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

**Hypoxic induction.** The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with MIGFR in hypoxic conditions (such as with 0.1% O<sub>2</sub>, 5% CO<sub>2</sub>, and balance N<sub>2</sub>, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses an MIGFR, and that optionally has defective IGFR function (e.g. IGFR is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate IGFR modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate IGFR modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether MIGFR function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express MIGFR relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the MIGFR plays a direct role in hypoxic induction.

**Cell adhesion.** Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., *Bioconjug Chem.* 2001 May-Jun;12(3):346-53).

**Tubulogenesis.** Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include Matrigel<sup>TM</sup> (Becton Dickinson), an extract of basement membrane proteins

containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4° C and forms a solid gel at 37° C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, *Nature Medicine* 5:1418-1423). These assays have traditionally involved stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF- $\alpha$ . Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing an MIGFR's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF- $\alpha$ , ephrin, etc.

**Cell Migration.** An invasion/migration assay (also called a migration assay) tests the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, *J Biol Chem* 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are

fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing an MIGFR's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

**Sprouting assay.** A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 $\mu$ l of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100  $\mu$ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

#### ***Primary assays for antibody modulators***

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the MIGFR protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for

detecting MIGFR-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

### ***Primary assays for nucleic acid modulators***

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance MIGFR gene expression, preferably mRNA expression. In general, expression analysis comprises comparing MIGFR expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express MIGFR) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that MIGFR mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, *eds.*, John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the MIGFR protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve MIGFR mRNA expression, may also be used to test nucleic acid modulators.

### **Secondary Assays**

Secondary assays may be used to further assess the activity of MIGFR-modulating agent identified by any of the above methods to confirm that the modulating agent affects MIGFR in a manner relevant to the IGFR pathway. As used herein, MIGFR-modulating agents encompass candidate clinical compounds or other agents derived from previously

identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with MIGFR.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express MIGFR) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate MIGFR–modulating agent results in changes in the IGFR pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use “sensitized genetic backgrounds”, which, as used herein, describe cells or animals engineered for altered expression of genes in the IGFR or interacting pathways.

#### ***Cell-based assays***

Cell based assays may detect endogenous IGFR pathway activity or may rely on recombinant expression of IGFR pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

#### ***Animal Assays***

A variety of non-human animal models of normal or defective IGFR pathway may be used to test candidate MIGFR modulators. Models for defective IGFR pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the IGFR pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, IGFR pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal IGFR are used to test the candidate modulator's affect on MIGFR in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at

4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the MIGFR. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on MIGFR is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, e.g., Ogawa K et al., 2000, *Oncogene* 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the MIGFR endogenously are injected in the flank,  $1 \times 10^5$  to  $1 \times 10^7$  cells per mouse in a volume of 100  $\mu$ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorigenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate

modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for evaluation of the candidate modulator.

Tumorigenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorigenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorigenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

#### **Diagnostic and therapeutic uses**

Specific MIGFR-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the IGFR pathway, such as angiogenic, apoptotic, or cell proliferation disorders.



Accordingly, the invention also provides methods for modulating the IGFR pathway in a cell, preferably a cell pre-determined to have defective or impaired IGFR function (e.g. due to overexpression, underexpression, or misexpression of IGFR, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates MIGFR activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the IGFR function is restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought closer to normal compared to untreated cells. For example, with restored IGFR function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired IGFR function by administering a therapeutically effective amount of an MIGFR -modulating agent that modulates the IGFR pathway. The invention further provides methods for modulating MIGFR function in a cell, preferably a cell pre-determined to have defective or impaired MIGFR function, by administering an MIGFR -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired MIGFR function by administering a therapeutically effective amount of an MIGFR -modulating agent.

The discovery that MIGFR is implicated in IGFR pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the IGFR pathway and for the identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether MIGFR expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (e.g., *Current Protocols in Molecular Biology* (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, *Biotechniques* (1999) 26:112-125; Kallioniemi OP, *Ann Med* 2001, 33:142-147; Blohm and Guiseppi-Elie, *Curr Opin Biotechnol* 2001, 12:41-47). Tissues having a disease or disorder implicating defective IGFR signaling that express an MIGFR, are identified as amenable to treatment with an MIGFR modulating agent. In a preferred application, the IGFR defective tissue overexpresses an MIGFR

relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial MIGFR cDNA sequences as probes, can determine whether particular tumors express or overexpress MIGFR. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of MIGFR expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the MIGFR oligonucleotides, and antibodies directed against an MIGFR, as described above for: (1) the detection of the presence of MIGFR gene mutations, or the detection of either over- or under-expression of MIGFR mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of MIGFR gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by MIGFR.

Kits for detecting expression of MIGFR in various samples, comprising at least one antibody specific to MIGFR, all reagents and/or devices suitable for the detection of antibodies, the immobilization of antibodies, and the like, and instructions for using such kits in diagnosis or therapy are also provided.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in MIGFR expression, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for MIGFR expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer. The probe may be either DNA or protein, including an antibody.

## **EXAMPLES**

The following experimental section and examples are offered by way of illustration and not by way of limitation.

### **I. *Drosophila* IGFR screen**

A dominant loss of function screen was carried out in *Drosophila* to identify genes that interact with or modulate the IGFR signaling pathway. Activation of the pathway by overexpression of IGFR at early stages in the developing *Drosophila* eye leads to an increase in cell number which results in a larger and rougher adult eye (Potter CJ et al. (2001) Cell 105:357-368; Huang et al., 1999. Dev. 126:5365-5372). We generated a fly stock with an enlarged eye due to overexpression of IGFR and identified modifiers of this phenotype. We then identified human orthologues of these modifiers.

The screening stock carried two transgenes. The genotype is as follows:

+; +; *P{DmIGFR-pExp-UAS}* *P{Gal4-pExp-1Xey}/TM6B*

Screening stock females of the above genotype were crossed to males from a collection of 3 classes of piggyBac-based transposons. The resulting progeny, which contain both the transgenes and the transposon, were scored for the effect of the transposon on the eye overgrowth phenotype (either enhancement, suppression or no effect). All data was recorded and all modifiers were retested with a repeat of the original cross.

## II. Analysis of Table 1

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of *Drosophila* modifiers. The columns “MIGFR symbol”, and “MIGFR name aliases” provide a symbol and the known name abbreviations for the Targets, where available, from Genbank. “MIGFR RefSeq\_NA or GI\_NA”, “MIGFR GI\_AA”, “MIGFR NAME”, and “MIGFR Description” provide the reference DNA sequences for the MIGFRs as available from National Center for Biology Information (NCBI), MIGFR protein Genbank identifier number (GI#), MIGFR name, and MIGFR description, all available from Genbank, respectively. The length of each amino acid is in the “MIGFR Protein Length” column.

Names and Protein sequences of *Drosophila* modifiers of IGFR from screen (Example I), are represented in the “Modifier Name” and “Modifier GI\_AA” column by GI#, respectively.

Table 1

MIGFR symbol	MIGFR name aliases	MIGFR RefSeq_NA or GI_NA	MIGFR GI_AA	MIGFR name	MIGFR description	MIGFR protein length	Modifier name	Modified gi_aa
NOS3	nitric oxide synthase 3 (endothelial cell) ECNOS eNOS NOS3	NM_00603	40254422	nitric oxide synthase 3 (endothelial cell)	nitric oxide synthase; electron transporter; apoptosis inhibitor	1203	Nos	24583543
PRKAG1	AMPK gamma 1 PRKAG1 Protein kinase, AMP-activated, noncatalytic, gamma 1 protein kinase, AMP-activated, gamma 1 non-catalytic subunit	NM_002733	4506061	protein kinase, AMP-activated, gamma 1 non-catalytic subunit	protein kinase; protein serine/threonine kinase; protein binding	331	SNF4 Agamma	24648661
PRKAG2	AAKG AAKG2 H91620P H91620p protein PRKAG2 WPWS protein kinase, AMP-activated, gamma 2 non-catalytic subunit	NM_016203	33186925	protein kinase, AMP-activated, gamma 2 non-catalytic subunit	SNF1A/AMP-activated protein kinase; AMP binding	569	SNF4 Agamma	24648661
PRKAG3	AMPK gamma 3 protein kinase, AMP-activated, gamma 3 non-catalytic subunit PRKAG3	NM_017431	8394044	protein kinase, AMP-activated, gamma 3 non-catalytic subunit	nucleotide binding; protein binding; SNF1A/AMP-activated protein kinase	492	SNF4 Agamma	24648661
NOS2A	HEP-NOS INOS NOS NOS, type II NOS2 NOS2A nitric oxide synthase 2A (inducible, hepatocytes) nitric oxide synthase, macrophage	NM_00625 NM_153292	24041029	nitric oxide synthase 2A (inducible, hepatocytes)	calcium channel inhibitor; nitric oxide synthase; calmodulin binding; electron transporter; antiviral response protein	1153	Nos	24583543
ITPKA	IP3K IP3K A IP3K-A ITPKA inositol 1,4,5-	NM_002220	4504789	inositol 1,4,5-trisphosphate 3-kinase A	1D-myo-inositol-trisphosphate 3-kinase	461	IP3K1	24583132

	ol 1,4,5-trisphosphate 3-kinase A			e 3-kinase A				
ITPKB	IP3 3-kinase IP3K IP3K B IP3KB ITPKB inositol 1,4,5-trisphosphate 3-kinase B	NM_002221	38569400	inositol 1,4,5-trisphosphate 3-kinase B	1D-myo-inositol-trisphosphate 3-kinase	946	IP3K1	24583132
ITPKC	IP3K C IP3K-C ITPKC inositol 1,4,5-trisphosphate 3-kinase C	NM_025194	18643383	inositol 1,4,5-trisphosphate 3-kinase C	inositol/phosphatidylinositol kinase	683	IP3K1	24583132
MAN2A1	Golgi alpha-mannosidase II mannosidase, alpha, class 2A, member 1 mannosidase, alpha type II MANA2 MAN2A1	NM_002372	4758698	mannosidase, alpha, class 2A, member 1	mannosyl-oligosaccharide 1,3-1,6-alpha-mannosidase	1143	alpha-Man-IIb	24647249
MAN2A2	alpha-mannosidase IIX mannosidase, alpha, class 2A, member 2 manosidase, alpha-, type II, isozyme X mannosidase, alpha type II-X PRO2198 MANA2X MAN2A2	NM_006122 NM_018621	3123244	mannosidase, alpha, class 2A, member 2	alpha-mannosidase	1139	alpha-Man-IIb	24647249
NOS1	nitric oxide synthase 1 (neuronal) NOS NOS1	NM_000620	10835173	nitric oxide synthase 1 (neuronal)	nitric oxide synthase; electron transporter	1434	Nos	24583543
KIAA1815	FLJ23309 KIAA1815	XM_291315	37540252	KIAA1815	na	904	CG11961	24655610
POMT1	POMT1 RT protein-O-mannosyltransferase 1	NM_007171	21361382	protein-O-mannosyltransferase 1	mannosyltransferase	747	rt	10727985

PTPRN	IA-2 IA-2 PTP IA2 ICA512 PTPRN R-PTP-N islet cell antigen 2 islet cell antigen 512 islet cell autoantigen 3 protein tyrosine phosphatase, receptor type, N protein tyrosine phosphatase-like N precursor	NM_002846	4506321	protein tyrosine phosphatase, receptor type, N	transmembrane receptor protein tyrosine phosphatase; transmembrane receptor protein tyrosine phosphatase	979	ia2	24580758
PTPRN2	IA-2beta IAR IAR2beta IAR receptor-like protein-tyrosine phosphatase ICAAAR KIAA0387 PHOGRIN PTPRN2 PTPRP phogrin protein tyrosine phosphatase receptor pi protein tyrosine phosphatase, receptor type, N polypeptide 2 tyrosine phosphatase IA-2 beta	NM_002847 NM_130842 NM_130843	11386149	protein tyrosine phosphatase, receptor type, N polypeptide 2	transmembrane receptor protein tyrosine phosphatase	1015	ia2	24580758
SYNJ1	INPP5G SYNJ1 inositol 5'-phosphatase (synaptojanin 1) synaptojanin 1 synaptojanin-1, polyphosphoinositide phosphatase	NM_003895	4507335	synaptojanin 1	polyphosphoinositide phosphatase; inositol-1,4,5-trisphosphate 5-phosphatase	1575	synaptojanin	18497296
SYNJ2	INPP5H SYNJ2 inositol phosphate 5'-phosphatase 2 (synaptojanin 2) synaptojanin 2	NM_003898	26190608	synaptojanin 2	protein binding; inositol/phosphatidylinositol phosphatase; inositol-1,4,5-trisphosphate 5-phosphatase	1496	synaptojanin	18497296

LOC284767	na similar to dJ680N4.2 (ubiquitin-conjugating enzyme E2D 3 (homologous to yeast UBC4/5)) LOC 284767	XM_209367	27485630	similar to dJ680N4.2 (ubiquitin-conjugating enzyme E2D 3 (homologous to yeast UBC4/5))	na	147	eff	24646906
LOC51619	ubiquitin-conjugating enzyme HBUCE1 LOC 51619	NM_015983	8393719	ubiquitin-conjugating enzyme HBUCE1	ubiquitin conjugating enzyme; enzyme	147	eff	24646906
UBE2D2	ubiquitin-conjugating enzyme E2D 2 (UBC4/5 homolog, yeast) ubiquitin-conjugating enzyme E2-17 kDa 2 ubiquitin carrier protein E2(17)KB2 UBCH5B UBC4/5 PUBC1 UBC4 UBE2D2	NM_003339 NM_181838	33188456	ubiquitin-conjugating enzyme E2D 2 (UBC4/5 homolog, yeast)	ubiquitin conjugating enzyme;	118	eff	24646906
UBE2D3	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast) ubiquitin-conjugating enzyme E2-17 kDa 3 ubiquitin carrier protein E2(17)KB3 MGC43926 MGC5416 UBCH5C UBC4/5 UBE2D3	NM_003340 NM_181886 NM_181887 NM_181888 NM_181889 NM_181890 NM_181891 NM_181892 NM_181893	33149322	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast)	ubiquitin conjugating enzyme	148	eff	24646906
LOC378372	LOC378372 na similar to dopamine beta-hydroxylase-like	XM_353788	37539365	similar to dopamine beta-hydroxylase-like	na	451	olf413	28574720
MOXD1	DKFZP564G202 MOX MOXD1 dJ248E1.1 monooxygenase, DBH-like 1	NM_015529	37183305	monooxygenase, DBH-like 1	dopamine-beta-monooxygenase; copper binding; monooxygenase	613	olf413	28574720

### III. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled MIGFR peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of MIGFR activity.

### IV. High-Throughput In Vitro Binding Assay.

<sup>33</sup>P-labeled MIGFR peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl<sub>2</sub>, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate IGFR modulating agents.

### V. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins,  $3 \times 10^6$  appropriate recombinant cells containing the MIGFR proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at  $15,000 \times g$  for 15 min. The cell lysate is incubated with 25  $\mu$ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel



electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

#### VI. Kinase assay

A purified or partially purified MIGFR is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10  $\mu\text{g/ml}$ ). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100  $\mu\text{l}$ . The reaction is initiated by the addition of  $^{33}\text{P}$ -gamma-ATP (0.5  $\mu\text{Ci/ml}$ ) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation ( $\text{Mg}^{2+}$  or  $\text{Mn}^{2+}$ ) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

#### VII. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues are obtained from Impath, UC Davis, Clontech, Stratagene, Ardaïs, Genome Collaborative, and Ambion.

TaqMan® analysis is used to assess expression levels of the disclosed genes in various samples.

RNA is extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/μl. Single stranded cDNA is then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan® assay (Applied Biosystems, Foster City, CA) are prepared according to the TaqMan® protocols, and the following criteria: a) primer pairs are designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis is performed using a 7900HT instrument.

TaqMan® reactions are carried out following manufacturer's protocols, in 25 μl total volume for 96-well plates and 10 μl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis is prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data are normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples are compared with matched normal tissues from the same patient. A gene is considered overexpressed in a tumor when the level of expression of the gene is 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue is not available, a universal pool of cDNA samples is used instead. In these cases, a gene is considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type is greater than 2 times the standard deviation of all normal samples (i.e.,  $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$  ).

A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression

of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

WHAT IS CLAIMED IS:

1. A method of identifying a candidate IGFR pathway modulating agent, said method comprising the steps of:
  - (a) providing an assay system comprising a MIGFR polypeptide or nucleic acid;
  - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
  - (c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as a candidate IGFR pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the MIGFR polypeptide.
3. The method of Claim 2 wherein the cultured cells additionally have defective IGFR function.
4. The method of Claim 1 wherein the assay system includes a screening assay comprising a MIGFR polypeptide, and the candidate test agent is a small molecule modulator.
5. The method of Claim 4 wherein the assay is a binding assay.
6. The method of Claim 1 wherein the assay system is selected from the group consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay comprising a MIGFR polypeptide and the candidate test agent is an antibody.

8. The method of Claim 1 wherein the assay system includes an expression assay comprising a MIGFR nucleic acid and the candidate test agent is a nucleic acid modulator.
9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
11. The method of Claim 1 additionally comprising:
  - (d) administering the candidate IGFR pathway modulating agent identified in (c) to a model system comprising cells defective in IGFR function and, detecting a phenotypic change in the model system that indicates that the IGFR function is restored.
12. The method of Claim 11 wherein the model system is a mouse model with defective IGFR function.
13. A method for modulating a IGFR pathway of a cell comprising contacting a cell defective in IGFR function with a candidate modulator that specifically binds to a MIGFR polypeptide, whereby IGFR function is restored.
14. The method of claim 13 wherein the candidate modulator is administered to a vertebrate animal predetermined to have a disease or disorder resulting from a defect in IGFR function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
16. The method of Claim 1, comprising the additional steps of:
  - (d) providing a secondary assay system comprising cultured cells or a non-human animal expressing MIGFR ,

(e) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and

(f) detecting an agent-biased activity of the second assay system,  
wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate IGFR pathway modulating agent,

and wherein the second assay detects an agent-biased change in the IGFR pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a IGFR pathway gene.

20. A method of modulating IGFR pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a MIGFR polypeptide or nucleic acid.

21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the IGFR pathway.

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

23. A method for diagnosing a disease in a patient comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the sample with a probe for MIGFR expression;
- (c) comparing results from step (b) with a control;

(d) determining whether step (c) indicates a likelihood of disease.

24. The method of claim 23 wherein said disease is cancer.

**ABSTRACT OF THE DISCLOSURE**

Human MIGFR genes are identified as modulators of the IGFR pathway, and thus are therapeutic targets for disorders associated with defective IGFR function. Methods for identifying modulators of IGFR, comprising screening for agents that modulate the activity of MIGFR are provided.



### Nucleic Acid and Polypeptide sequences

```
>gi|40254421|ref|NM_000603.2| Homo sapiens nitric oxide synthase 3
(endothelial cell) (NOS3), mRNA
AAACTAGGGGCAAGGAGACGAAGAGAACATGAAAGTTAAACTTTAAGATGAAGAACAAGCTGAACATAC
TGATGCATTGGATCTTTGGAGAGGATCTCAGAATCATTGTACTTAATTTACAGGCTAAAACCTTAGAAG
AGGAATTTATTATATCCACACAAGACTCCAGGGAAGCACATGGCCCTGGACTGAAGGCTGGCATCTGGA
AGCTGTTCAGCCACCAGCACCTTCTGCAGCAGGAAAAGGCCAGGGCTCTGCTGGAGCAGGCAGCAGAGTGG
ACGCACAGTAACATGGGCAACTTGAAGAGCGTGGCCCAGGAGCCTGGGCCACCCTGCGGCCCTGGGGCTGG
GGCTGGGCCCTTGGGCTGTGCGGCAAGCAGGGCCCAGCCACCCCGGCCCTGAGCCCAGCCGGGCCCCAGC
ATCCCTACTCCACCAGCGCCAGAACACAGCCCCCGAGCTCCCCGCTAACCAGCCCCAGAGGGGGCCC
AAGTTCCCTCGTGTGAAGAACTGGGAGGTGGGGAGCATCACCTATGACACCCTCAGCGCCCAGGCGCAGC
AGGATGGGCCCTGCACCCCAAGACGCTGCCTGGGCTCCCTGGTATTTCCACGGAACTACAGGGCCGGCC
CTCCCCCGGCCCCCGGCCCTGAGCAGCTGCTGAGTCAGGCCCCGGGACTTCATCAACCAGTACTACAGC
TCCATTAAGAGGAGCGGCTCCAGGCCCACGAACAGCGGCTTCAAGAGGTGGAAGCCGAGGTGGCAGCCA
CAGGCACCTACCAGCTTAGGGAGAGCGAGCTGGTGTTCGGGGCTAAGCAGGCC TGGCGCAACGCTCCCCG
CTGCGTGGGGCCGATCCAGTGGGGGAAGCTGCAGGTGTTGATGCCC GGGACTGCAGGTCTGCACAGGAA
ATGTTACCTTACATCTGCAACCACATCAAGTATGCCACCAACCGGGGCAACCTTCGCTCGGCCATCACAG
TGTTCCCGCAGCGCTGCCCTGGCCGAGGAGACTTCCGAATCTGGAACAGCCAGCTGGTGC GCTACGCGGG
CTACCGGCAGCAGGATGGCTCTGTGCGGGGGGACCCAGCCAACGTGGAGATCACCGAGCTCTGCATT CAG
CACGGCTGGACCCCAGGAAACGGTCGCTTCGACGTGCTGCCCTTGCTGCTGCAGGCCCCAGATGATCCCC
CAGAATCTTTCCTTCTGCCCCCGAGCTGGTCCCTTGAGGTGCCCC TGGAGCACCCACGCTGGAGTGGTT
TGCAGCCCTGGGCC TGCCTGGTACGCCCTCCCGGCAGTGTCCAACATGCTGCTGGAATTTGGGGGCTG
GAGTTCCCGCAGCCCCCTTCACTGGCTGGTACATGAGCACTGAGATCGGCACGAGGAACCTGTGTGACC
CTACCGCTACAACATCTGGAGGATGTGGCTGTCTGCATGGACCTGGATACCCGACCACCTCGTCCCT
GTGGAAGACAAGGCAGCAGTGGAAATCAACGTGGCCGTGCTGCACAGTTACAGCTAGCCAAAGTCACC
ATCGTGGCAACACAGCGCCACGGCTCTTTTCATGAAGCACTGGAGAATGAGCAGAAGGCCAGGGGGG
GCTGCCCTGCAGACTGGGCC TGGATCGTGCCCCCTCATCGGGCAGCCTCAC TCCTGTTTTCCATCAGGA
GATGGTCAACTATTTCTGTCCCCGGCCTTCCGCTACAGCCAGACCCCTGGAAGGGGAGTGC CGCCAAG
GGCACCGCATCACAGGAAGAAGACCTTTAAAGAAGTGGCCAACGCCGTGAAGATCTCCGCC TCGCTCA
TGGGCACGGTGTATGGCGAAGCAGTGAAGGCGACAATCCTGTATGGCTCCGAGACCGGCCGGGCCAGAG
CTACGCACAGCAGCTGGGGAGACTCTTCCGGAAGGCTTTTGATCCCCGGGTCTGTGTATGGATGAGTAT
GACGTGGTGTCCCTCGAACACGAGACGCTGGTGTGGTGGTAACCAGCACATTTGGGAATGGGGATCCCC
CGGAGAATGGAGAGAGCTTTGCAGCTGCCC TGATGGAGATGTCCGGCCCCCTACAACAGCTCCCC TCGGCC
GGAACAGCACAAAGAGTTATAAGATCCGCTTCAACAGCATCTCCTGCTCAGACCCACTGGTGTCTCTTTGG
CGGCGGAAGAGGAAGGAGTCCAGTAACACAGACAGTGCAGGGGGCCCTGGGCACCC TCAGGTTCTGTGTGT
TCGGGCTCGGCTCCCGGGCATACCCCACTTCTGCGCCTTTGCTCGTGCCGTGGACACACGGCTGGAGGA
ACTGGGCGGGGAGCGGCTGCTGCAGCTGGGCCAGGGCGACGAGCTGTGCGGCCAGGAGGAGGCCTTCCGA
GGCTGGGGCCAGGCTGCCCTTCAGGCCGCCTGTGAGACCTTCTGTGTGGGAGAGGATGCCAAGGCCGCCG
CCCCAGACATCTTCAGCCCCAAACGGAGCTGGAAGCGCCAGAGGTACCGGCTGAGCGCCAGGCCGAGGG
CCTGCAGTTGCTGCCAGGTCTGATCCACGTGCACAGGCGGAAGATGTTCCAGGCTACAATCCGCTCAGTG
GAAAACCTGCAAAGCAGCAAGTCCACGAGGGCCACCATCCTGGTGCGCC TGGACACCGGAGGCCAGGAGG
GGCTGCAGTACCAGCCGGGGGACCACATAGGTGTCTGCCCGCCAACCGGCCGGCCTTGTGGAGGCGCT
GCTGAGCCGCGTGGAGGACCCGCCGGCGCCCACTGAGCCCGTGGCAGTAGAGCAGCTGGAGAAGGGCAGC
CCTGGTGGCCCTCCCCCGGCTGGGTGCGGGACCCCGGCTGCCCGCTGCACGCTGCGCCAGGCTCTCA
CCTTCTTCTTGGACATCACCTCCCCACCCAGCCCTCAGCTCTTGCGGCTGCTCAGCACCTTGGCAGAAGA
GCCCAGGGAACAGCAGGAGCTGGAGGCCCTCAGCCAGGATCCCCGACGCTACGAGGAGTGAAGTGGTTTC
CGCTGCCCCACGCTGCTGGAGGTGCTGGAGCAGTTCCCGTGGTGGCGCTGCCTGCCCCACTGCTCCTCA
CCCAGCTGCCCTCTGCTCCAGCCCCGGTACTACTCAGTCAGCTCGGCACCCAGCACCCACCCAGGAGAGAT
CCACCTCACTGTAGCTGTGCTGGCATAACAGGACTCAGGATGGGCTGGGCCCCCTGCATATGGAGTCTGC
TCCACGTGGCTAAGCCAGCTCAAGCCGGAGACCCGTGTGCCCTGCTTCATCCGGGGGGCTCCCTCCTTCC
GGCTGCCACCCGATCCCAGCTTGCCCTGCATCCTGGTGGGTCCAGGCACTGGCATTGCCCCCTTCCGGGG
ATTCTGGCAGGAGCGGCTGCATGACATTGAGAGCAAAGGGCTGCAGCCCACTCCCATGACTTTGGTGTTC
GGCTGCCGATGCTCCCAACTTGACCATCTCTACCGCAGCAGGTGCAGAACGCCAGCAGCGCGGGGTGT
TTGGCCGAGTCTCACCGCTTCTCCCGGGAACCTGACAACCCCAAGACCTACGTGCAGGACATCCTGAG
GACGGAGCTGGCTGCGGAGGTGCACCGCGTGTGTGCCCTCAGCGGGGCCACATGTTTGTCTGCGGCGAT
GTTACCATGGCAACCAACGTCTGTCAGACCGTGCAGCGCATCTGGCGACGGAGGGCGACATGGAGCTGG
```

ACGAGGCCGGCGACGTATCGGGCGTGCTGCGGGATCAGCAACGCTACCACGAAGACATTTTCGGGGCTCAC  
 GCTGCGCAGCCAGGAGGTGACAAGCCGCATACGCACCCAGAGCTTTTTCCTTGACAGGAGCGTCAGTTGCGG  
 GGCGCAGTGCCCTGGGGCGTTTCGACCCCTCCCGGCTCAGACACCAACAGCCCCGTGAGAGCCGCTGGCTTTT  
 CCTTCCAGTTCCGGGAGAGCGGCTGCCCCGACTCAGGTCCGCCCCGACCAGGATCAGCCCCGCTCCTCCCCCT  
 CTTGAGGTGGTGCCTTCTCACATCTGTCCAGAGGCTGCAAGGATTCAGCATTTATTCCTCCAGGAAGGAGC  
 AAAACGCCTCTTTTCCCTCTCTAGGCCTGTTGCCTCGGGCCTGGGTCCGCTTAATCTGGAAGGCCCTC  
 CCAGCAGCGGTACCCAGGGCCTACTGCCACCCGCTTCTGTTTCTTAGTCGAATGTTAGATTCTCTTG  
 CCTCTCTCAGGAGTATCTTACCTGTAAAGTCTAATCTCTAAATCAAGTATTTATTATTGAAGATTTACCA  
 TAAGGGACTGTGCCAGATGTTAGGAGAACTACTAAAGTGCTTACCCAGCCCCAAAAAAAAAAAAAAAAAAAA

>gi|4506060|ref|NM\_002733.1| Homo sapiens protein kinase, AMP-  
 activated, gamma 1 non-catalytic subunit (PRKAG1), mRNA

GCGCCCTTAAAGATGGTGAGGGGGCTCATGCTCTGAGTAGAAGGTGGTGACCTCCAGGAGCGGTGGGATG  
 ATGAGGGCCCCGGGCGCCTCTTGCAATGGAGACGGTCATTTCTTCAGATAGCTCCCCAGCTGTGGAAAATG  
 AGCATCCTCAAGAGACCCAGAAATCCAACAATAGCGTGTATACTTCCTTCATGAAGTCTCATCGCTGCTA  
 TGACCTGATTCCCACAAGCTCCAAATTGGTTGTATTTGATACGTCCCTGCAGGTGAAGAAAGCTTTTTTTT  
 GCTTTGGTGACTAACGGTGTACGAGCTGCCCCCTTTATGGGATAGTAAGAAGCAAAGTTTTGTGGGCATGC  
 TGACCATCACTGATTTTCATCAATATCCTGCACCGCTACTATAAATCAGCCTTGGTACAGATCTATGAGCT  
 AGAAGAACAAGATAGAACTTGGAGAGAGGTGTATCTCCAGGACTCCTTTAAACCGCTTGTCTGCATT  
 TCTCCTAATGCCAGCTTGTGGTGTCTCTCTTCAATTAATTCGGAACAAGATCCACAGGCTGCCAGTTA  
 TTGACCCAGAATCAGGCAATACTTTGTACATCCTCACCCACAAGCGCATCTGAAGTTCTCAAATTGTT  
 TATCACTGAGTTCCCCAAGCCAGAGTTCATGTCCAAGTCTCTGGAAGAGCTACAGATTGGCACCTATGCC  
 AATATTGCTATGGTTTCGCACTACCACCCCCGCTCTATGTGGCTCTGGGGATTTTTGTACAGCATCGAGTCT  
 CAGCCCTGCCAGTGGTGGATGAGAAGGGGGCGTGTGGTGGACATCTACTCCAAGTTTGATGTTATCAATCT  
 GGCAGCAGAAAAGACCTACAACAACCTAGATGTATCTGTGACTAAAGCCTTGCAACATCGATCACATTAC  
 TTTGAGGGTGTCTCAAGTGCTACCTGCATGAGACTCTGGAGACCATCATCAACAGGCTAGTGGAAGCAG  
 AGGTTACCCGACTTGTAGTGGTGGATGAAAATGATGTGGTCAAGGGAATTGTACTGTCTGACATCCT  
 GCAGGCCCTGGTGTCTCAGGTGGAGAGAAGAAGCCCTGAGCTGGGGGAAGGGGTCTGCAGCACCAGGG  
 GATATGCCCAACTCACTGCCGTCTGGAAGCTCTGTGGGAATCAGATGAACTTGAGGGAATTGTGACTCT  
 GTTCCCTGTTTCCAGGTCCCCGCCCCCTTCTATCTGGGAGCTAGGGAAGGTATGGGGGAGGAAAGAGCAATGG  
 ATTTATAGCTATCCCTTACCCCTCACACATACACTTGAAAAAACTTTTCCAGCTAGCCAGTTAGACCCCTGT  
 CCTCTTAGATATATCCCCCTTTCTGGGTGAACACTATAGGCTCTGTGCCCTCTCAGACAAATCTGATCTCTA  
 AGAGATCCCCAGACCTCACTTGCCTCTGCCTCCATCTTGGCCCTGATTCAACCCCTAAGATAATAGCACAA  
 CAAAATTCTTCATAAAGATATTTTTATTACCTGTTCCGTGCTATATGGAGGAGGCCAAGTCCATTTAGT  
 GACATTTCTTCCATAATGTGAGTGGGGAGGATTGTGG

>gi|33186924|ref|NM\_016203.2| Homo sapiens protein kinase, AMP-  
 activated, gamma 2 non-catalytic subunit (PRKAG2), mRNA

GCGGCTCCAGTCCGAAGGCAGCGCCGGGGGAGGGAAGGAGGGGACCGAACCCCGAGGAGTTTCGCAGA  
 ATCAACTTCTGGTTAGAGTTATGGGAAGCGCGGTTATGGACACCAAGAAGAAAAAAGATGTTTCCAGCCC  
 CGGCGGGAGCGGCGGCAAGAAAAATGCCAGCCAGAAGAGGCGTTTCGCTGCGCGTGCACATTCCGGACCTG  
 AGCTCCTTCGCCATGCCGCTCCTGGACGGAGACCTGGAGGGTTCGCGAAAGCATTCCTCTCGAAAGGTGG  
 ACAGCCCCCTTCGGCCCGGGCAGCCCCCTCCAAAGGGTTCTTCTCAGAGGCCCCCAGCCCCGGCCCTCCAG  
 CCCCATGTCTGCACCTGTGAGGCCCAAGACCAGCCCCGGCTCTCCCAAACCGTGTTCCTGTTCTCTTAC  
 CAGGAGTCCCGCCACGCTCCCCTCGACGCATGAGCTTCAGTGGGATCTTCCGCTCCTCCTCCAAAGAGT  
 CTTCCCCCAACTCCAACCTTGCTACCTCGCCCGGGGCATCAGGTTTTCTCCCGCTCCAGAAAAACCTC  
 CGGCCTCTCCTCCTCTCCGTCAACACCCACCCAAGTGACCAAGCAGCACACGTTTTCCCTGGAACTCTAT  
 AAGCACGAGCCTGAACGGTTAGAGAATCGCATCTATGCCTCGTCTTCCCCCGGACACAGGGCAGAGGT  
 TCTGCCCGTCTTCTTCCAGAGCCCGACAGGCCCTCCACTGGCATCACCGACACACTATGCTCCCTCCAA  
 AGCCGCGGCGCTGGCGGGCGGCCCTGGGACCCCGGAAGCCGGCATGTGGAGAAGCTGGAGTTCGAGGAC  
 GAAGCAGTAGAAGACTAGAAAGTGGTGTACATGCGATTTCATGAGGTACACAAGTGTTATGACATCG  
 TTCCAACAGTTCAAAGCTTGTTGTCTTTGATACTACATTACAAGTTAAAAAGGCCTTCTTTGCTTTGGT  
 AGCCAACGGTGTCCGAGCAGCGCCACTGTGGGAGAGTAAAAAACAAAGTTTTGTAGGAATGCTAACAATT  
 ACAGATTTTATAAATATACTACATAGATACTATAAATCACCTATGGTACAGATTTATGAATTAGAGGAAC  
 ATAAAATTGAAACATGGAGGGAGCTTTATTTACAAGAAACATTTAAGCCTTTAGTGAATATATCTCCAGA  
 TGCAAGCCTCTTCGATGCTGTATACTCCTTGATCAAAAATAAAATCCACAGATTGCCCGTTATTGACCT  
 ATCAGTGGGAATGCATTTTATATACTTACCCACAAAAGAATCCTCAAGTTCTCAGCTTTTTATGTCTG

ATATGCCAAAGCCTGCTTTTCATGAAGCAGAACCTGGATGAGCTTGGAAATAGGAACGTACCACAACATTGC  
 CTTTCATACATCCAGACACTCCCATCATCAAAGCCTTGAACATATTTGTGGAAAGACGAATATCAGCTCTG  
 CCTGTTGTGGATGAGTCAGGAAAAGTTGTAGATATTTATTCCAAATTTGATGTAATTAATCTTGCTGCTG  
 AGAAAACATACAATAACCTAGATATCACGGTGACCCAGGCCCTTCAGCACCGTTTCACAGTATTTTGAAGG  
 TGTTGTGAAGTGCAATAAGCTGGAAATACTGGAGACCATCGTGGACAGAATAGTAAGAGCTGAGGTCCAT  
 CGGTTGGTGGTGGTAAATGAAGCAGATAGTATTGTGGGTATTATTTCCCTGTCCGACATTCTGCAAGCCC  
 TGATCCTCACACCAGCAGGTGCCAAACAAAAGGAGACAGAAACGGAGTGACCGCCGTGAATGTAGACGCC  
 CTAGGAGGAGAACTTGAACAAAGTCTCTGGGTCACGTTTTGCCTCATGAACACTGGCTGCAAGTGGTTAA  
 GAATGTATATCAGGGTTTAAACGATAGGTATTTCTTCCAGTGATGTTGAAATTAAGCTTAAAAAAGAAAGA  
 TTTTATGTGCTTGAATAATTCAGGCTTGCATTAATAAACTGTTTTTCAGACCTTGTCTGAAGGATTTTAAAT  
 GCTGTATGTCTATTAAAGTGCACGTGTCTCCCTG

>gi|8394043|ref|NM\_017431.1| Homo sapiens protein kinase, AMP-  
 activated, gamma 3 non-catalytic subunit (PRKAG3), mRNA

ACTTGGTCTGGGGCTGGCCACATGGAGCCCGGGCTGGAGCACGCACTGCGCAGGACCCCTTCCTGGAGCA  
 GCCTTGGGGGTTC TGAGCATCAAGAGATGAGCTTCCTAGAGCAAGAAAACAGCAGCTCATGGCCATCACC  
 AGCTGTGACCAGCAGCTCAGAAAGAATCCGTGGGAAACGGAGGGCCAAAGCCTTGAGATGGACAAGGCAG  
 AAGTCGGTGGAGGAAGGGGAGCCACCAGGTGAGGGGAAGGTCCCCGGTCCAGGCCAGCTGCTGAGTCCA  
 CCGGGCTGGAGGCCACATTCCCCAAGACCACACCCTTGGCTCAAGCTGATCC TGCCGGGGTGGGCACTCC  
 ACCAACAGGGTGGGACTGCCCTCCCC TCTGACTGTACAGCCTCAGCTGCAGGCTCCAGCACAGATGATGTG  
 GAGCTGGCCACGGAGTTCCAGCCACAGAGGCCTGGGAGTGTGAGCTAGAAGGCCTGCTGGAAGAGAGGC  
 CTGCCCTGTGCCCTGTCCCCGCAGGCCCATTTCCCAAGCTGGGCTGGGATGACGAAC TGCGGAAACCCGG  
 CGCCCAGATCTACATGCGCTTCATCGAGGAGCACACCTGCTACGATGCCATGGCAACTAGCTCCAAGCTA  
 GTCATCTTCGACACCATGCTGGAGATCAAGAAGGCCTTCTTTGCTCTGGTGGCCAACGGTGTGCGGGCAG  
 CCCCTCTATGGGACAGCAAGAAGCAGAGCTTTGTGGGGATGCTGACCATCACTGACTTCATCTCTGGTGT  
 GCATCGCTACTACAGGTCCCCCTTGGTCCAGATCTATGAGATTGAACAACATAAGATTGAGACCTGGAGG  
 GAGATCTACCTGCAAGGTGCTTCAAGCCTCTGGTCTCCATCTCTCC TAATGATAGCCTGTTTGAAGCTG  
 TCTACACCCTCATCAAGAACCGGATCCATCGCCTGCCTGTTCTTGACCCGGTGTGAGGCAACGTACTCCA  
 CATCTCACACACAAACGCCCTGCTCAAGTTCTTGCACATCTTTGGTTCCCTGCTGCCCCGGCCCCCTCTTC  
 CTCTACCGCACTATCCAAGATTTGGGCATCGGCACATTCCGAGACTTGGCTGTGGTGTGAGAGACAGCAC  
 CCATCCTGACTGCACATGGACATCTTTGTGGACCGGCGTGTGCTGCACTGCCCTGTGGTCAACGAATGTGG  
 TCAGGTGCTGGGGCTCTATTCCCGCTTTGATGTGATTCACCTGGCTGCCCAGCAAACCTACAACCACCTG  
 GACATGAGTGTGGGAGAAGCCCTGAGGAAGAGGACACTATGTCTGGAGGGAGTCCTTTCTGCCAGCCCC  
 ACGAGAGCTTGGGGGAAGTGATCGACAGGATTGCTCGGGAGCAGGTACACAGGCTGGTGTAGTGGACGA  
 GACCCAGCATCTCTTGGGCGTGGTCTCCCTCTCCGACATCTTCAGGCACTGGTGTGCTAGCCCTGCTGGC  
 ATCGATCCCTCGGGGCC TGAGAAGATCTGAGTCCCTCAATCCCAAGCCACCTGCACACCTGGAAGCCAATG  
 AAGGGAAC TGGAGAAC TCAAGCTTCACTCTTCCCCCACCCTTCTGCTGGTTCAGCTATGATTGAGGCTT  
 CTTCAGGCCCC TCCAAATTGCCCTTGCCCTACCAGAGCTCCCAAGCCCTCGGCATGCCAGTGCACCAT  
 GGGATGATGAAATTAAGGAGAACAGCTGAGTCAAGCTTGGAGGTCCCTGAACCAGAGGCACTAGGATTAC  
 CCCAGGGCCATCTGTGCTCCATGCCCCCCCATCCCCCTTGCCGCTGACTGGGTGCGATGGCCCCCAGTGG  
 GTTTAGTCAGGGCTTCTGGATCTCGGTTTCTGGGCTACCTATGCTTCACCTTCAGCTCCTGGGAGTTCC  
 CAGCAACGTGCGCCACTGCCCCCTCTACTCTCCAGGGGTTGGTCATTTCAAGGCTGCTGAAATGCTGCATT  
 TCAGGGGGCCACCATGGAGCAGCCGTTATTTATAGAAGTGCCCTGTTGGAGGTGGGGAGTCTCCCTCCA  
 TTCTTGTCCAGAAAACCTCTTAGCTCTCGCAGTGAGCCCATGTTCTTAGTCTCCAGGGATGGATGGCCTT  
 GTATATGGACCCCTGAGAATGAGCAATTGAGAAAACAAAACAAAAGGAACAATCCATGAACTTAGATTTT  
 ATTGGTTTCACTCAAAATGCTGCAGTCATTTGACCTGAACTTGTGGCAAGAGACTTGTGCTTTCTAAATT  
 CAAAGACTAGAAGGGAATGGATAAAAATCACAAGTGCCGTTTCTCTTG

>gi|24041028|ref|NM\_000625.3| Homo sapiens nitric oxide synthase 2A  
 (inducible, hepatocytes) (NOS2A), transcript variant 1, mRNA

ATAACTTTGTAGCGTAGCGAACTGAGGCTCCGGCCGAGAGAACTCAGCCTCATTCCTGCTTTAAAT  
 CTC TCGGCCACCTTTGATGAGGGGACTGGGCAGTTCTAGACAGTCCCGAAGTTCTCAAGGCACAGGTCTC  
 TTCTGGTTTACTGTCTTTACCCCGGGGAGGCAGTGACCCAGCTGCAAGCCCCACAGTGAAGAACATC  
 TGAGCTCAAATCCAGATAAGTGACATAAGTGACCTGCTTTGTAAAGCCATAGAGATGGCCTGTCTTGGA  
 AATTTCTGTTCAAGACCAAATTCCACCAGTATGCAATGAATGGGGAAAAAGACATCAACAACAATGTGGA  
 GAAAGCCCCCTGTGCCACCTCCAGTCCAGTGACACAGGATGACCTTCAGTATCACAACCTCAGCAAGCAG  
 CAGAATGAGTCCCCGAGCCCCCTCGTGGAGACGGGAAAGAAGTCTCCAGAATCTCTGGTCAAGCTGGATG

CAACCCATTGTCCTCCCCACGGCATGTGAGGATCAAAAACCTGGGGCAGCGGGATGACTTTCCAAGACAC  
 ACTTCACCATAAGGCCAAAGGGATTTTAACTTGCAGGTCCAAATCTTGCCTGGGGTCCATTATGACTCCC  
 AAAAGTTTGACCAGAGGACCCAGGGACAAGCCTACCCCTCCAGATGAGCTTCTACCTCAAGCTATCGAAT  
 TTGTCAACCAATATTACGGCTCCTTCAAAGAGGCAAAAATAGAGGAACATCTGGCCAGGGTGGGAAGCGGT  
 AACAAAGGAGATAGAAACAACAGGAACCTACCAACTGACGGGAGATGAGCTCATCTTCGCCACCAAGCAG  
 GCCTGGCGCAATGCCCCACGCTGCATTGGGAGGATCCAGTGGTCCAACCTGCAGGTCTTCGATGCCCCGA  
 GCTGTTCCACTGCCCCGGAAATGTTTGAACACATCTGCAGACACGTGCGTTACTCCACCAACAATGGCAA  
 CATCAGGTGCGCCATCACCGTGTCCCCAGCGGAGTGATGGCAAGCACGACTTCCGGGTGTGGAATGCT  
 CAGCTCATCCGCTATGCTGGCTACCAGATGCCAGATGGCAGCATCAGAGGGGACCTTGCCAACGTGGAAT  
 TCACTCAGCTGTGCATCGACCTGGGCTGGAAGCCCAAGTACGGCCGCTTCGATGTGGTCCCCCTGGTCT  
 GCAGGCCAATGGCCGTGACCTGAGCTCTTTCGAAATCCACCTGACCTTGTGCTTGAGGTGGCCATGGAA  
 CATCCCCAATACGAGTGGTTCGGGAACCTGGAGCTAAAGTGGTACGCCCTGCCTGCAGTGGCCAACATGC  
 TGCTTGAGGTGGGCGGCTGGAGTTCACAGGTGCCCTTCAATGGCTGGTACATGGGCACAGAGATCGG  
 AGTCCGGGACTTCTGTGACGTCCAGCGCTACAACATCCTGGAGGAAGTGGGCAGGAGAATGGGCCTGGAA  
 ACGCACAAGCTGGCCTCGCTCTGGAAAGACCAGGCTGTGCTTGAGATCAACATTGCTGTGCTCCATAGTT  
 TCCAGAAGCAGAATGTGACCATCATGGACCACCACTCGGCTGCAGAATCCTTCATGAAGTACATGCAGAA  
 TGAATACCGGTCCCGTGGGGGCTGCCCGGCAGACTGGATTGGCTGGTCCCTCCCATGTCTGGGAGCATC  
 ACCCCCGTGTTCACCAGGAGATGCTGAACTACGTCTGTCCCTTTCTACTACTATCAGGTAGAGGCCT  
 GGAAAACCCATGTCTGGCAGGACGAGAAGCGGAGACCCAAGAGAAGAGAGATTCCATTGAAAGTCTTGGT  
 CAAAGCTGTGCTCTTTGCCTGTATGCTGATGCGCAAGACAATGGCGTCCCGAGTCAGAGTCACCATCCTC  
 TTTGCGACAGAGACAGGAAAATCAGAGGCCTGGCCTGGGACCTGGGGGCCCTTATTCAGCTGTGCCTTCA  
 ACCCCAAGGTTGTCTGCATGGATAAGTACAGGCTGAGCTGCCTGGAGGAGGAACGGCTGCTGTTGGTGGT  
 GACCAGTACGTTTGGCAATGGAGACTGCCCTGGCAATGGAGAGAACTGAAGAAATCGCTCTTCATGCTG  
 AAAGAGCTCAACAACAAATTCAGGTACGCTGTGTTTGGCCTCGGCTCCAGCATGTACCCTCGGTTCTGCG  
 CCTTTGCTCATGACATTGATCAGAAGCTGTCCCACCTGGGGGCCCTCAGCTCACCCCGATGGGAGAAGG  
 GGATGAGCTCAGTGGGCAGGAGGACGCCTTCCGCAGCTGGGCCGTGCAAACCTTCAAGGCAGCCTGTGAG  
 ACGTTTGATGTCCGAGGCAACAGCACATTACAGATCCCCAAGCTCTACACCTCCAATGTGACCTGGGACC  
 CGCACCACTACAGGCTCGTGCAGGACTCACAGCCTTGGACCTCAGCAAAGCCCTCAGCAGCATGCATGC  
 CAAGAACGTGTTACCATGAGGCTCAAATCTCGGCAGAATCTACAAAGTCCGACATCCAGCCGTGCCACC  
 ATCCTGGTGGAACTCTCCTGTGAGGATGGCCAAGGCCTGAAC TACCTGCCGGGGGAGCACCTTGGGGTTT  
 GCCCAGGCAACCAGCCGGCCCTGGTCCAAGGTATCTGGAGCGAGTGGTGGATGGCCCCACACCACCA  
 GACACTGAGCCCTGAGGCCCCCTGGATGAGAGTGGCAGCTACTGGGTGAGTACAAGAGGCTGCCCCCTGC  
 TCACTCAGCAGGCCCCCTACCTACTTCCCTGGACATCACACACCCCCAACCCAGCTGCTGCTCCAAAAGC  
 TGGCCCAGGTGGCCACAGAAGAGCCTGAGAGACAGAGGCTGGAGGCCCTGTGCCAGCCCTCAGAGTACAG  
 CAAGTGGAAAGTTACCAACAGCCCCACATTCTTGGAGGTGCTAGAGGAGTTCCCGTCCCTGCGGGTGTCT  
 GCTGGCTTCCCTGCTTTCCAGCTCCTCCATTCTGAAGCCCAGGTTCTACTCCATCAGCTCCTCCCGGATC  
 ACACGCCCACAGAGATCCACCTGACTGTGGCCGTGGTCACTACCACACCCGAGATGGCCAGGGTCCCT  
 GCACCACGGCGTCTGCAGCACATGGCTCAACAGCCTGAAGCCCCAAGACCCAGTGGCCTGCTTTGTGCGG  
 AATGCCAGCGGCTTCCACCTCCCCGAGGATCCCTCCCATCCTTGCATCCTCATCGGGCTGGCACAGGCA  
 TCGCGCCCTTCCGAGTTTCTGGCAGCAACGGCTCCATGACTCCCAGCACAAAGGAGTGCGGGGAGGCCG  
 CATGACCTTGGTGTGTTGGGTGCCGCCGCCAGATGAGGACCACATCTACCAGGAGGAGATGCTGGAGATG  
 GCCCAGAAGGGGTGCTGCATGCGGTGCACACAGCCTATTCCCGCCTGCCTGGCAAGCCCAAGGTCTATG  
 TTCAGGACATCCTGCGGCAGCAGCTGGCCAGCGAGGTGCTCCGTGTGCTCCACAAGGAGCCAGGCCACCT  
 CTATGTTTGGGGGATGTGCGCATGGCCCCGGGACGTGGCCCCACACCCTGAAGCAGCTGGTGGCTGCCAAG  
 CTGAAATTGAATGAGGAGCAGGTCGAGGACTATTTCTTTTCAAGCTCAAGAGCCAGAAGCGCTATCACGAAG  
 ATATCTTTGGTGTGTATTTCTTACGAGGCGAAGAAGGACAGGGTGGCGGTGCAGCCCAGCAGCCTGGA  
 GATGTGACGCTCTGAGGGCTACAGGAGGGGTAAAGCTGCCGGCACAGAACTTAAGGATGGAGCCAGC  
 TCTGCATTATCTGAGGTACAGGGCCTGGGGAGATGGAGGAAAGTGATATCCCCAGCCTCAAGTCTTAT  
 TTCTCAACGTTGCTCCCCATCAAGCCCTTTACTTGACCTCCTAACAAGTAGCACCCCTGGATTGATCGGA  
 GCCTCCTCTCTCAAACCTGGGGCCTCCCTGGTCCCTTGGAGACAAAATCTTAAATGCCAGGCCCTGGCAAGT  
 GGGTGAAGATGGAACCTGCTGCTGAGTGACCACTTCAAGTGACCACAGGAGGTGCTATCGCACCACT  
 GTGTATTTAACTGCCTTGTGTACAGTTATTTATGCCTCTGTATTTAAAAAACTAACACCCAGTCTGTTCC  
 CCATGGCCACTTGGGTCTTCCCTGTATGATTCTTGGATGGAGATATTTACATGAATTGCATTTTACTTTA  
 ATCACAAAAAAAAAAAAAAAAAA

>gi|4504788|ref|NM\_002220.1| Homo sapiens inositol 1,4,5-trisphosphate 3-kinase A (ITPKA), mRNA

GAATTCCGGAAATGACCTGCCCGGGGGCCCCAACGGGCATGGCGCGGCCGGGGGGCGCAGGGCCCTGCAG  
 CCCGGGGCTGGAGCGGGCCCCGCGCCGGAGTGTCTGGGGAGCTGCGCCTGCTCTTCGAGGCGCGCTGTGCG  
 GCGGTTCGTGCGGGCCGCCGCGCGGGGGAGCCCCGGGCCCCGCGGGGCCAAGCGGCCTGGGGGACAGGTCC  
 CCAACGGGCTTCCGCGGGCTCCCCCGGGCCCCGGTGATCCCTCAGCTGACCGTGACAGCCGAGGAGCCCGA  
 CGTGCCCCCGACCAGCCCTGGGCGCGCGGAGCGGGAGAGGGACTGCCCTCCCGGCAGCGGGCTCTTCGCAC  
 CTGCAGCAGCCGCGCCGCCCTTTCACCTCGTCGGTCTCTCTCCACTGGCTCCTCGTCGCTGCTCGAGGACT  
 CGGAGGACGACCTGCTGAGCGACAGTGAGAGCCGGAGCCGCGGCAACGTGCAGCTGGAAGCGGGCGAGGA  
 CGTGGGTGAGAAAACCACTGGCAGAAGATCCGGACCATGGTCAATCTGCCGGTCATAAGCCCTTTCAAG  
 AAGCGTACGCCCTGGGTGACGTGGCAGGGCACACTGGGAGTTTAAAGCGCGCGGGCACCAGCGGGCTGA  
 TCCTGAAGCGCTGCTCGGAGCCGGAGCGCTACTGCCCTGGCGCGGCTGATGGCTGACGCGCTGCGCGGCTG  
 CGTGCTTGCCTTTCACGGCGTGGTGGAGCGCGACGGCGAAAGCTACCTGCAGCTGCAGGACCTGCTCGAT  
 GGCTTCGACGGACCTTGTGTGCTCGACTGCAAAAATGGGCGTCAAGGACTTACCTAGAGGAGGAGCTGACCA  
 AGGCCCCGTGAGCGGCCAAGCTGCGGAAGGACATGTACAAGAAAATGCTGGCGGTGGATCCTGAAGCTCC  
 CACGGAGGAGGAGCAGCGCAGCGCGCCGTCACCAAGCCGCGCTACATGCAGTGGCGGGAAGGCATCAGC  
 TCCAGCACCCCTCGGCTTCCGCATCGAGGGCATCAAGAAAGCGGACGGCTCCTGCAGCACCGACTTCA  
 AGACTACGCGAAGCCGAGAGCAGGTGCTTCGCGTCTTTGAAGAGTTTGTGCAAGGAGATGAGGAAGTGCT  
 GAGGCGGTATCTGAACCGCTGACGACAGATCCGGGACACCCCTGGAGGTATCCGAGTTCTTCAGGAGGCAC  
 GAGGTGATCGGCAGCTCGCTCCTCTTTGTGACGATCACTGCCATCGCGCCGGCGTGTGGCTCATCGACT  
 TCGGCAAGACCACGCCCCCTCCCGATGGCCAGATCTTGGACCACCGGCGGCCCTGGGAGGAGGGCAACCG  
 CGAGGACGGCTATTTGCTGGGGCTGGACAATCTCATTTGGCATCTTGGCCAGCCTGGCTGAGAGATGAGGC  
 TGGACTCCTGTCCCCGCGGGCCGCTCACCTGACATGTGGACCTGCAGCTTTGTCCCACTGTGCATGCCG  
 GCTTGAGACTGGAGCCCCGCGGTGCAGGGCAGTTACCGGGTCTTGCAGGACCAGGTGCCAGCCACTAAG  
 GGGGGGCACCGCCGATGCCAGGGGTTTTGCCACCCGGGCCCCAGCGTTCCAGAGCCAAATGACACTAA  
 CTTATAGAAGGGGAGGGGGCAAAGGGCTTCTTCTCAGGCCAGCTCTTCTGAGGAGGCTCTGCCCTCTCC  
 AGAGGTGCCAGACCGCGGATTTTATTTAGCAAGCCAGACCTTCCGGTCTAACGTCTCACACCACGACGG  
 ACTCCCCCTTCTAATAAACTCAAAGACAAAA

>gi|38569399|ref|NM\_002221.2| Homo sapiens inositol 1,4,5-trisphosphate  
 3-kinase B (ITPKB), mRNA

GGAGCCGCGGCGGGCGGGCAGCGCGGGACCCAGTACTATGGCTGTGTACTGCTATGCGCTCAATAGCCTGG  
 TGATCATGAATAGCGCCAACGAGATGAAGAGCGGCGGGCGGCCCGGGGCCAGTGGCAGCGAGACGCCCC  
 GCCCCAGGAGGGCAGTGCTGAGCCCCGCGAGCGTTTTTCAGCCCCGGGAGAGGCGCCTCTTTCCTCTTC  
 CCCCCAGCCGAGTGTCTCCCCGAGGAGCCCCGAGCCCCGGGGGCTGGCGGAGCGGCCGGCGCAGGC  
 TGAATAGTAGCAGCGGCAGTGGCAGCGGCAGCAGCGGCAGTAGCGTGAGCAGCCCAAGTTGGGCTGGTCG  
 CCTGCGAGGGGACCGGCAGCAGGTGGTGGCAGCCGGTACCCCTCTCCCCGCCAGGGCCGGAGGAGGCCAAG  
 AGGAAGCTGCGGATCTTGCAGCGCGAGTTGCAGAACGTGCAGGTGAACCAGAAAAGTGGGCATGTTTGAAG  
 CGCACATCCAGGCACAGAGCTCCGCCATTCAAGCGCCCCGCAGCCCGCGTTTGGGCAGGGCTCGCTCGCC  
 CTCCCCGTGCCCCCTCCGCGAGCAGCAGTCAGCCCCCTGGAAGGGTCTTGGTTTCAAGGGCGCCGGAGCGAG  
 GAACGGAGGACAAAGTCTTGGGGGAGCAATGTCCAGAGACTTCAGGAACCGACTCCGGGAGGAAAGGAG  
 GGCCCAGCCTATGCTCTCGCAGGTGAAGAAAGGAATGCCACCCTTCTCCCGGCCGGGCTGCCCCCTACAGG  
 ATCAGAGGCTCAGGGTCCATCCGCTTTTGTAAAGGATGGAGAAGGGTATCCCTGCCAGTCCCCGCTGTGGC  
 TCACCCACAGCTATGGAAATTGACAAAAGGGGCTCTCCTACCCCGGGAACCTGGAGCTGCCTAGCTCCCT  
 CATTGGGGCTGTTCGGAGCTAGCTTAACGATGGCCACGGAAGTGGCAGCGAGAGTTACATCCACTGGGCC  
 ACACCGTCCACAGGATCTTGCCCTCACTGAGCCGTCTGGGAGAGCCCGTGAGCTTGAGGACCTGCAGCCC  
 CCAGAGGCCCCCTGGTGGAGAGGCAGGGGCAGTTTCTGGGCAGTGAGACAAGCCCAGCCCCAGAAAGGGGCG  
 GGCCCCGCGATGGAGAACCCCCCTGGGAAGATGGGGAAGGATATCTGCCCTGTGGCATGCCGGGCTCTGG  
 GGAGCCTGAAGTGGGCAAAAGGCCAGAGGAGACGACTGTGAGCGTGCAAAGCGCAGAGTCTCTGATTCC  
 CTGAGCTGGTCCAGGCTGCCAGGGCCCCCTGGCCTCCGTAGGCCCTGAGGAGGCCCGAAGTGGGGCCCCCG  
 TGGGCGGGGGGCGTTGGCAGCTCTCCGACAGAGTGGAGGGAGGGTCCCCAACGCTGGGCTTGCTTGGGGG  
 CAGCCCCCTGACACAGCCGGGGACCGGGAATGTGGAGGCGGGAATTCCTTTCGGCAGAATGCTGGAGCCT  
 TTGCCCTGTGTGGGACGCTGCGAAAGATCTGAAAGAAGCTCAGTGCCCTCTTGGGGACAGGGTGGGTGTC  
 AGCCTGGGAACCTCAGGGTTTGGCAGGGCACCATGGAGAAAGCCGGTTTGGCTTGGACGCGTGGCACAGG  
 GGTGCAATCAGAGGGGACTTGGGAAAGCCAGCGGCAGGACAGTGATGCCCTCCCAAGTCCGGAGCTGCTA  
 CCCCCAAGATCAGGACAAGCCCTTCTCTGAGGAAGGCTGCAGC CCCAGCAACATACCTGCTGTTCATCATTA  
 CAGACATGGGCACCCAGGAGGATGGGGCTTGGAGGAGACGCAGGGAAGCCCTCGGGGCAACCTGCCCTT  
 GAGGAAACTGTCTCTTCTCGGCCCTCTCCACGGGCTTCTCTCATCTCTACGAAGACTCAGAGGAGGAC  
 ATCTCCAGTGACCCTGAGCGCACCTGGACCCCAACTCAGCCTTCTTGCATACCTGGACCAGCAGAAAC

CTAGAGTGAGCAAATCATGGAGGAAGATAAAAAACATGGTGCACTGGTCTCCCTTCGTTCATGTCCCTTCAA  
 GAAGAAGTACCCCTGGATCCAGCTGGCAGGACACGCAGGGAGTTTCAAGGCAGCTGCCAATGGCAGGATC  
 CTGAAGAAGCACTGTGTAGTCAGAGCAGCGCTGCCTGGACCGGCTGATGGTGGATGTGCTGAGGCCCTTCG  
 TACCTGCCTACCATGGGGATGTGGTGAAGGACGGGGAGCGCTACAACCAGATGGACGACCTGCTGGCCGA  
 CTTTCGACTCGCCCTGTGTGATGGACTGCAAGATGGGAATCAGGACCTACCTGGAGGAGGAGCTCACGAAG  
 GCGCGGAAGAAGCCCAGCCTGCGGAAGGACATGTACCAGAAGATGATCGAGGTGGACCCGAGGCCCCCA  
 CCGAGGAGGAAAAAGCACAGCGGGCTGTGACCAAGCCACGGTACATGCAGTGGCGGGAGACCATCAGCTC  
 CACGGCCACCCTGGGGTTTCAGGATCGAGGGAATCAAGAAAGAAGACGGCACCGTGAACCGGGACTTCAAG  
 AAGACCAAAACGAGGGAGCAGGTACCGAGGCCCTTCAGAGAGTTCACTAAAGGAAACCATAACATCCTGA  
 TCGCCTATCGGGACCGGCTGAAGGCCATTGCAACCACCTAGAAAGTTTCTCCCTTCTTCAAGTGCCACGA  
 GGTCAATTGGCAGCTCCCTCCTTTCATCCACGACAAGAAGAAACAGGCCAAAGTGTGGATGATCGACTTT  
 GGGAAAACCAGCCCCCTGCCTGAGGGCCAGACCCTGCAGCATGACGTCCCTGGCAGGAGGGGAACCGGG  
 AGGATGGCTACCTCTCGGGGCTCAATAACCTCGTCGACATCCTGACCGAGATGTCCCAGGATGCCCCACT  
 CGCCTGAGCTGCCCACGCCCTCCCTGGCCCCCGCTGGGCCTCCTTTCTCCTCCTGTGCTTCCTTTCTC  
 GTTCTTAACCTTTTCTTCACTTACACCTGACTGACCCTCCTGAACTGCACTACAAGACACTTTGTAGAAG  
 AGGAGATGAGAGTTTCTAGTCATTTTCTTAACCTTCAGGGCTTGAGGTGGTGTGTTGCACTGCTTTTGTGA  
 GAGAGGTACCTACTAGAAGAGAAATGCCAGTCTTAGAGGTGGGTGAGGTGTAGAGCTGGAGGGGGTTC  
 CCTGGCTGCTGAGGGGACCCTACCAGATGAGCCCTGCCTCTGGGAGCCCCCTAGGAAGCACCAGCCTGGA  
 CCTACCACCTGCGGAGGCTGCTGCCCCCTGGCGGCCAGTGCTGTTAGAGTGCTGCCAAGCACAGCCTTA  
 TTTCTGCGGGGGCTCCCCACCGAGAGCCAGGGGGCCGGCCGGTTCTGGTCCCTGGCTGGGAGCAG  
 GGCTTTCTGGTAGTTGGGGCACAAACCATCGGGGAACCATGTTGACTGTGAGCAAAGTGTCTTCCGA  
 TTAGCAGCCTCAGGGATGCCCTGGTGGCTCTCCAGGGCTGCTCAGGCAAGGCCCCCACCATCTGGTA  
 TGGAAACCTGCCGGCTCAGGCCAGACCCAGGAGCCAAGAGAAGGCTGAAGCCAGCTTGGCTGTGTTCTC  
 TGATCTAGGCCTTCCCAGAGGAGGCGAGCAGAAGCTGTGCCACTTGGAATTGCAACCCATGAGTTCAGAA  
 GGCACACTCTGCCATGCTGAGCTCCAAGGGTGTACCAGGGGAAGATGGGATCTATAGAGTCTCTGGGCC  
 CTGGCCCCAGGGAGGAGCACATTTTCTTGACCCTCACCTACCTGGTGTAGTTGGTCAACCCTGCCTGC  
 ATACATGGGCTCCTGTTCATGGGGCCAGAGTCCCTTGACAGATATAGAAATAGGGGAGGAGCTCAGGTCTG  
 CGCCAGGCAGGAAGAAGGCAGGCTTCTGGCTTCCAGAGGTGCCGCGGTGGCTCCTGGCATCATTTGTTA  
 TTGCCCTTGAAACAAGCCTTACTGCCCTGGAGGGCTTAGATTCCTGCTTCTCCAATGTAGTGTGGGTATCT  
 TGTAGGGTATGTGGTGGATGCCAGGGCGTGTCCAGGCACCTCTTCTGAAAGTCTCTGCATTTGGAGATT  
 CGTGAGAACCTATTTAAGCCCAATTTTAACAGAAAGCAAGTGTGATATGGAAGGAAATGTAATAAT  
 TTGCCTGACTTCTTAAGAACAACACCCAGCTCTGTGCCCCATGCTCCTTGGGGCTTGCCACCCACTCC  
 TTTGCTGTTCAGAGGTACAGGAGCTGGGAGAGTCCAGGAGCTAGGGACACAGAGGGAGACTATGGACCAAG  
 GTGTGTGTGCTGAGGAACCACTGCCCCACCCACCCACCCCGGGGTCCTGAGGAACTGTCAACCTGCC  
 ACGGGACATGTACATTTCCCTTTTGTGCTGGAAGTGTGAGTGACACTTGTGAGGGGTGGAGGGTGGGAC  
 ACATGAGGATGTATAAGTACAGATTTTAAAAAGGAAATCAACTTACACTTCTGGCTCTTGTTTAAAC  
 AGTGGTGAGCTCCTGTGTGGGCCGACTTGTCTAAAGGTACACACGCGCCGGTGGAGCACGAGAGACCTC  
 GTGGCAGCATGTGATCTGGAAGGCAGGCAGGACGGGGCGTTGGGGAGCCAAAGTCAACTCTGGGCCTCT  
 GGAGCTATAGTGACTTTTGGGCTAGAAGGGACCTGGTGGTCTGTGCTTTCAGCCATTTGCAGGGCAGGGG  
 CATCATTAATTCAGACGTAAAGATTCTATGAATATGGACTGGCCAAAAGTTATCCTTACTCCATCTGTGA  
 AAGAAGTTTGTCTAAAGCAAATCATGATATGAACAAAATTACAGGGGACCTGTTTAAGAGAAACAAATGT  
 TCCAAGCACTTTAGGCAGACACCAGCTGTTTGAACAATGTGCTAATATGCAAATGATGTGCTTATTAA  
 AGGAGGCCCATGGGGCTCTTATTGGCAATACCTGGCTGTGGGTACATTAAATATGTGAACATAGTATG  
 AAGTAGCATCATTTTAGGGTTATTCTGTTACTTAGGGTTTTTGTCTTCTGTTTTTTTTTCTCTTTTTTT  
 GTATTTACCGTGCTAGTTCTCTTCTACACCTACTCTGTCTCTCAAGCCATTTTGCCACTCGCTTCCCTGC  
 CATCTGGCCCTTCCCTTTGTCTCAGTGGGATAGATGGATTGTGAAATGGAATCTCCAGAACCCCTGCCC  
 TGGCAGCCTGGAAGACCGTGCTGCCCAGCCCTCGTCACCACAGGGACTCCTTGGGTCTGGCAGTGCAT  
 GTGCCAGCAGGCAGGACAACTCTGTGTACCTGTGCCCAGGTGAATGGGCGCAGGGTCTCTTTGCCCTGT  
 CCTGCGGGGGGCCCCACGAGTTCTTGGCATTACAGACTGCTTAGCATCTCGGAAGGTTTCTTCAAGTGC  
 TTGCTTTTCCAGGCTTGCCTTTAGTGTCTATGAAGACTTTTTTAAGTTATATTTATTTTGGGTTTTT  
 AAAATTGCACGAACACTAAGACCGAAAGGCTGGACTCTTGTCTTCTCCTTGAAAGCTTTGCCCTTTGTTTT  
 GAACCTTCTTTCCACTTGGTAGAAAGAGCCAGAAGCAGCCCTGGCCCTGTAAGATGGACTCTTTTCATC  
 CTTTCAGTTGTATTTAGCTTTGAGTTTCTCTGCATCTGTCCACCCCATGTGTATATAACCCAGCCCCCTGGC  
 TCTGGGGTGGTCACCTCGTCAGTGCCTTTTTGTCTGGAGGAGAGGACCCCCCGCCTGCCGAGAGGCTC  
 TCTTCTGTCTGACCCCTCTCCCCATGGGACCTTGGAGAAAAGTGAAGTTTACAAACCCCTGCACAG  
 TGCCTGTCAAACAGATGCAAACCTTCTGAAATAAGCCTTGGAGACCAAAAAAAAAAAAAAAAAAAAA

>gi|40254974|ref|NM\_025194.2| Homo sapiens inositol 1,4,5-trisphosphate 3-kinase C (ITPKC), mRNA

```
GGGTCCGGCCGAAGCCCGAAGGAGCGGGCATGAGGCGCTGCCCCGTGCCGTGGGAGCCTGAACGAGG
CGGAGGCCGGGGCGCTGCCCCGCGCGGGCCCGCATGGGACTGGAGGCGCCGCGAGGAGGGCGGGCGGCA
GCCGGGACAGCAGCGACCTGGGCCCGGCGCAGGGGCCCCGGCGGGGCGGCCGGAGGGGGCGGGCCCTGG
GCCCCGACAGAGGGGTCCAGCCTCCACAGCGAGCCTGAGAGGGCCGGCCTCGGGCCTGCGCCGGGGACAG
AGAGTCCGCAGGCAGAATTCCTGGACAGACGGACAGACTGAGCCCCGCGCAGCTGGCCTTGGAGTAGAGAC
CGAGAGGCCCAAGCAAAAGACGGAGCCAGACAGGTCCAGCCTCCGACGCATCTAGAATGGAGCTGGTCA
GAGCTGGAGACGACTTGTCTTTGGACGGAGACCGGGACAGATGGCCTTTGGACTGATCCGCACAGGTCCG
ACCTCCAGTTTCAGCCCCGAGGAGGCCAGCCCCCTGGACACAGCCAGGGGTTTCATGGGCCCTGGACAGAGCT
GGAAACGCATGGGTTCAGACTCAGCCAGAGAGGGTCAAGTCTGGGCTGATAACCTCTGGACCCACCAG
AACAGTTCCAGCCTCCAGACTCACCCAGAAGGAGCCTGTCCCTCAAAAGAGCCAAGTGCTGATGGCTCCT
GGAAAGAATTGTATACTGATGGCTCCAGGACACAACAGGATATTGAAGGTCCCTGGACAGAGCCATATAC
TGATGGCTCCAGAAAAAACAGGATACTGAAGCAGCCAGGAAACAGCCTGGCACTGGTGGTTTCCAAATA
CAACAGGATACTGATGGCTCCTGGACACAACCTAGCACTGACGGTTCCCAGACAGCACCTGGGACAGACT
GCCTCTTGGGAGAGCCTGAGGATGGCCCATTAGAGGAACCAGAGCCTGGAGAATTGCTGACTCACCTGTA
CTCTCACCTGAAGTGTAGCCCCCTGTGCCCTGTGCCCGCCTCATCATTACCCCTGAGACCCCTGAGCCT
GAGGCCAGCCAGTGGGACCCCCCTCCCGGGTTGAGGGGGGCGAGCGCGGCTTCTCCTCTGCCTCTTCTT
TCGACGAGTCTGAGGATGACGTGGTGGCCGGGGGCGGAGGTGCCAGCGATCCCAGGACAGGTCTGGGAG
CAAACCTTGAAGAAGCTGAAGACAGTTCTGAAGTATTCACCCCTTTGTGGTCTCCTTCCGAAACACTAC
CCTTGGGTCCAGCTTTCTGGACATGCTGGGAACCTCCAGGCAGGAGAGGATGGTCCGATTCTGAAACGTT
TCTGTCAGTGTGAGCAGCGCAGCCTGGAGCAGCTGATGAAAGACCCGCTGCGACCTTTCTGCTGCCTGCCTA
CTATGGCATGGTGTGTCAGGATGGCCAGACCTTCAACCAGATGGAAGACCTCCTGGCTGACTTTGAGGGC
CCCTCCATTATGGAAGTGAAGATGGGCAGCAGGACCTATCTGGAAGAGGAGCTAGTGAAGGCACGGGAAC
GTCCCCGTCCCCGGAAGGACATGTATGAGAAGATGGTGGCTGTGGACCCTGGGGCCCCCTACCCCTGAGGA
GCATGCCCAGGGTGCAGTCACCAAGCCCGCTACATGCAGTGGAGGGAACCATGAGCTCCACCTCTACC
CTGGGCTTCCGGATCGAGGGCATCAAGAAGGCAGATGGGACCTGTAACACCAACTTCAAGAAGACGCAGG
CACTGGAGCAGGTGACAAAAGTGTGGAGGACTTCGTGGATGGAGACCACGTCATCCTGCAAAAGTACGT
GGCATGCCTAGAAGAACTTCGTGAAGCTCTGGAGATCTCCCCCTTCTTCAAGACCCACGAGGTGGTAGGC
AGCTCCCCTCCTCTTCGTGCACGACCACACCGGCCGGCCAAAGGTCTGGATGATAGACTTCGGCAAGACGG
TGGCCTTGCCCCGACCACAGACGCTCAGCCACAGGCTGCCCTGGGCTGAGGGCAACCTGAGGACGGCTA
CCTCTGGGGCTGGACAACATGATCTGCCTCCTGCAGGGGCTGGCACAGAGCTGAGCTGCTCAGCCACCA
TCAGGTTAATTGGATGGCGCCAGTCTGGCTGGAGGAGCCCTGAGATGCCATGGGAGGCCTGAGGTTGGCC
ACGGGGGAGCTGGCCTCCAGGGACGGGAGAGATTGTGTTCATGTGCCACACGAGACCAACGTGGAAGAGTC
TGAAGGGCCTTGGGAGACCAGGTAGCACCTGGCCCCATCATGATGCAGGGGTTTTGGGGACCTGGAAGGA
AGGTGATGAGGCAGTGAGTCAGAAAAACAGAACGGGGTCCCCGGATCTGCCGGGAAGGCTTCTGAGGGG
CTGCCCTGAGAGCATTACGTTACATGTACAGGGTATGGTGTGACAGGGTGCCTGTGGACACATGAATC
ACTTCTAACCTGCCTCCCTGTGAGCCTCCAGGCTGCCAGCTGGCTGAGGCCAGGGACTGGGTGAGGCTCA
TCTGTGGCGCCTCAGAGGGTCAGCATCATTGGTGAACAGATGCAGGCGCTGCTGGACCATCTGGGGAGAG
TGACAGTCCATGTCTTACCAGGGAGCCATTTGAGTGTGAGCGACAAGAGGCTCAGAGGGCATGACCCC
ATGGGACTGGATGCGGCCTGAGGGCTGATACCGCTGGGCGGTATCCTGCCCTGCTGTGGCCCTGTGGGAT
CCTCCGTGTTTCTCGGCGGACTCTGCTGACCTCCTGCAGACCCAAACCACAGCCACATCCCAGCTTCTGT
GCCAGCACTGTGACAGTACCTCGCTCCTCTGTGCACAGATCCGGCCTCAGGACTTACACCTCCTGCCTG
ACCCCCAGGCTTCTCTCTCCTTTCTCCCAGCAAACCTGCAGTGGCAGAAAGGAGGTTTCAAGGCTGGGAAA
GTGGGCCTCCCCCTTGCAACTCAGAGCTGCTGCACTCAGGAGGGCCCCATCCAATCCCGGGCCCCCTGCAGG
GAAAAGCGCTGGGTGTGTGTCAGAGGCGCAGGGTGGGTGGGGCTGCCAGCCAGGACCCCTGGCCTGCAGCC
TGATCCAAACCAAAGACTGTAGAACCCTGGGGTGTGGCTAACGGCCCCCTCCAGCACCCATAGCCAGGTCT
TCCTGGCCCTTGGAGCTGGGCTGGCGGACAGGCACCTACCTCTTCCCTTAAGCTGAAGCTCCACACTGTC
TTCCAGGGCTGAGGAGATGCTCTCCTTTTCTACTGACCATCTTGATACTTATTTATACGAGAGGCAGTTG
CTGGACGGGGTAGTACTGGGAAGCAGGAGGCAGAATGGCTCTGCTGAGCCTCCTACCCATGACAACACCC
CAATAAACAGAACATTCAGAGCCAAAAA
```

>gi|4758697|ref|NM\_002372.1| Homo sapiens mannosidase, alpha, class 2A, member 1 (MAN2A1), mRNA

```
GGCTCCTCGCGATCTTGTTCCTTTCCCCCTCCGCTTCTCTGACCTAGCTGCGCGGCCCCGGCCCGGGAGCT
GCCGAACCCGCGCCTCCCCCTGGGTGAGGAGGACACGCCTGCCCTCGTCGAGAAAACCTTTTCTGCCGACT
CAGTTGGGGCGGCGGTGGCAGGAAGTGCGGGCAGCGACCTCTCCTCCGCTGCCCGCGCGCCCTGCCGG
```

AGGTGCGGCTGAGCTTGCATCAAGTTTGTGGGGCCCCCTTCCCAGTTGCCGGCGAGTCTCGCCTCG  
 AGAGGGGGCGCCGACCCCGGGGAGGGCGGCAGGCCAGGGCGAAGGCCAAGGGCGTTTTGTGGCGCCGGAG  
 ACTAGGTGCGGAGCAAGGCGGGGACTCGCACCCGCATCCGAGAGCGCGGAGGTTCGCGCAGCCCCGGGAGAA  
 GGGAGCCTCCGGCGGGCTGCTTCTTAGAGTCCACAGTGCCTGTCTCCTTTGGCTGAGGAGAGTGTCTTGG  
 CCCCAGTCTATCGAGGAAAATGAAGTTAAGCCGCCAGTTTACCCTGTTTCGGCAGTGCATCTTCTGTGT  
 GGTGATTTTCTCGCTCTACCTGATGCTGGACCGGGGTCACTTAGACTACCCAGGAACCCGCGCCGCGAG  
 GGCTCCTTCCCTCAGGGCCAGCTCTCAATGTTGCAAGAAAAATAGACCATTTGGAGCGTTTGCTAGCTG  
 AGAATAATGAGATCATCTCAAATATTAGAGACTCAGTCATCAATTTGAGTGAGTCTGTGGAGGATGGTCC  
 GAAAAGTTACAAAAGCAATTTACGCCAAGGTGCTGGCTCACATCTTCTGCCCTCACAAATTATCCCTCTCA  
 GTTGACACTGCAGACTGTCTGTTTGTCTTCAAAAGTGAAGTACAATTAGATGTGCAGATGTTGGATG  
 TTTACAGTCTAATTTCTTTTGACAATCCAGATGGTGGAGTTTGAAGCAAGGATTTGACATTACTTATGA  
 ATCTAATGAATGGGACTGAACCCCTTCAAGCTTTGTGGTGCCTCATTCCCATAACGACCCAGGTTGG  
 TTGAAGACTTTCAATGACTACTTTAGAGACAAGACTCAGTATATTTTTAATAACATGGTCTTAAAGCTGA  
 AAGAAGACTCACGGAGGAAGTTTATTTGGTCTGAGATCTTTACCTTTCAAAGTGGTGGGATATTATAGA  
 TATTACAGAAGAAGGATGCTGTTAAAAGTTTAAATAGAAAATGGTCAGCTTGAAATTGTGACAGGTGGCTGG  
 GTTATGCCTGATGAAGCTACTCCACATTATTTTGCCTTAATTGATCAACTAATTGAAGGACATCAGTGGC  
 TGGAAAATAATATAGGAGTGAAACCTCGGTCCGGCTGGGCTATTGATCCCTTTGGACACTCACCAACAAT  
 GGCTTATCTTCTAAACCGTGTGGACTTTCTCACATGCTTATCCAGAGAGTTTATTATGCAGTTAAAAAA  
 CACTTTGCACTGCATAAAACATTGGAGTTTTTTTGGAGACAGAATTGGGATCTGGGATCTGTCACAGATA  
 TTTTATGCCACATGATGCCCTTCTACAGCTATGACATCCCTCACACTTGTGGACCTGATCCTAAAATATG  
 CTGCCAGTTTGATTTTAAACGTCTTCTGGAGGCAGATTTGGTTGTCCCTGGGGAGTCCCCCAGAAACA  
 ATACATCCTGGAAATGTCCAAAGCAGGGCTCGGATGCTACTAGATCAGTACCGAAAGAAGTCAAAGCTTT  
 TTCGTACCAAAGTTCTCCTGGCTCCACTAGGAGATGATTTCCGCTACTGTGAATACACGGAATGGGATTT  
 ACAGTTTAAAGATTATCAGCAGCTTTTTTGATTATATGAATTTCTCAGTCCAAGTTTAAAGTTAAGATACAG  
 TTTTGAACTTTATCAGATTTTTTTGATGCGCTGGATAAAGCAGATGAAACTCAGAGAGACAAGGGCCAGT  
 CGATGTTCCCTGTTTTAAGTGGAGATTTTTTCACTTATGCCGATCGAGATGATCATTACTGGAGTGGCTA  
 TTTTACATCCAGACCCTTTTACAAACGAATGGACAGAATCATGGAATCTCATTTAAGGGCTGCTGAAATT  
 CTTTACTATTTTCGCCCTGAGACAAGCTCACAAATACAAGATAAAATAAATTTCTCTCATCATCACTTTACA  
 CGGCACTGACAGAAGCCAGAAGGAATTTGGGACTGTTTCAACATCATGATGCTATCACAGGAAGTGCAAA  
 AGACTGGGTGGTGTGGATTATGGTACCAGACTTTTTCATTCTGTTAATGGTTTTGGAGAAGATAAATGGA  
 AATTCTGCATTTCTTCTTATTTTGAAGGACAACTCACATACGACTCTTACTCTCTGATACCTTCTCTGG  
 AGATGGATTGTAACAAAATACAAGATTCTCTGCCACAAAAAATATAATAAGGCTGAGTGCAGGAGCC  
 AAGGTACCTTGTGGTCTATAATCCTTTAGAACAAGACCGAATCTCGTTGGTCTCAGTCTATGTGAGTTCC  
 CCGACAGTGCAAGTGTCTCTGCTTTCAGGAAAACCTGTGGAAGTTCAAGTCAGCGCAGTTTGGGATACAG  
 CAAATACTATTTTCAAGAACAGCCTATGAGATCTCTTTTCGAGCACATATACCGCCATTGGGACTGAAAGT  
 GTATAAGATTTTGAATCAGCAAGTTCAAATTCACATTTAGCTGATTATGTCTTGTATAAGAATAAAGTA  
 GAAGATAGCGGAATTTTACCATAAAGAATATGATAAATACTGAAGAAGGTATAACACTAGAGAACTCCT  
 TTGTTTTACTTCGGTTTGTATCAAACGGACTTATGAAGCAAATGATGACTAAAGAAGATGGTAAACACCA  
 TGAAGTAAATGTGCAATTTTTCATGGTATGGAACCAATTTAAAGAGACAAAAGTGGTGCCTACCTCTTC  
 TTACCTGATGGTAATGCCAAGCCTTATGTTTACACAACACCGCCCTTTGTCAGAGTGACACATGGAAGGA  
 TTTATTCGGAAGTGACTTGCTTTTTTGACCATGTTACTCATAGAGTCCGACTATACCACATACAGGGAAT  
 AGAAGGACAGTCTGTGGAAGTTTCCAATATTGTGGACATCCGAAAAGTATATAACCGTGAGATTGCAATG  
 AAAATTTCTTCTGATATAAAAAGCCAAAATAGATTTTATACTGACCTAAATGGGTACCAGATTCAACCTA  
 GAATGACACTGAGCAAATTTGCCTCTTCAAGCAAATGTCTATCCCATGACCACAATGGCCTATATCCAGGA  
 TGCCAAACATCGTTTTGACACTGCTCTCTGCTCAGTCTTTAGGGGTTTCGAGTTTGAATAGTGGTCAGATT  
 GAAGTTATCATGGATCGAAGACTCATGCAAGATGATAATCGTGGCCTTGAGCAAGGTATCCAGGATAACA  
 AGATTACAGCTAATCTATTTTGAATACTACTAGAAAAAAGAAGTGTGTTAATACGGAAGAAGAAAAGAA  
 GTCGGTCAGTTATCTTCTCTCTTACCCACATAAATCTTCTCTCATGAATCATCCAGTCATTCCAATG  
 GCAAATAAGTTCTCACCTACCTTTGAGCTGCAAGGTGAATTCCTTCCATTACAGTCATCTTTGCCCTGTG  
 ACATTTCATCTGGTTAATTTGAGAACAATACAGTCAAAGGTGGGCAATGGGCACTCCAATGAGGACGCTT  
 GATCTTCAACAGAAAAAGGTTTGTGTCGTTCTCTAGCAAAGGCACAGGGCTGTTTTGTTCTACTACT  
 CAGGGAAAGATATTTGGTACAGAACTTTTAAACAAGTTTATTGTGCAAGTCTCACACCTTCATCACTAT  
 CCTTGATGCATTACCTCCCGGCACTCAGAATATAAGTGAGATCAACTTGAGTCCAATGGAAATCAGCAC  
 ATTCCGAATCCAGTTGAGGTGAACCTGACTTTTACATTTGGATTGAGAATCATTGGCTTTTATACCTTTC  
 TTGGTTTTCAGCTGCAATAAAGAAGCACATTATTTTAGCTTCTGGCTACTGTGAGAACATGAATTCTGTGA  
 TTCTGTGGGTTTTTTCTTTTTTCTTTTACCAGTACAGTAAG



>gi|5540099|ref|NM\_006122.1| Homo sapiens mannosidase, alpha, class 2A, member 2 (MAN2A2), mRNA

GGCAGCTCGGCCGACTGGGCCCCGAGCGGCGCGGAGGCCGGGCGCTGACGGTGTGTGTGGAGGCCAGTAT  
GAAGCTGAAAAAGCAGGTGACAGTGTGTGGGGCTGCCATCTTCTGTGTGGCAGTCTTCTCGCTCTACCTC  
ATGCTGGACCGAGTGAACACGATCCCACCCGACACCAGAATGGTGGGAAC'TCCCCCGAGCCAAATTT  
CTGTGCTGCAGAACCGCATTGAGCAGCTGGAGCAGCTTTTGGAGGAGAACCATGAGATTATCAGCCATAT  
CAAGGACTCCGTGCTGGAGCTGACAGCCAACGCAGAGGGCCCCGCCCATGCTGCCCTACTACACGGTC  
AATGGCTCCTGGGTGGTGCCACCGGAGCCCCGGCCAGCTTCTTCTCCATCTCCCCGAGGACTGCCAGT  
TTGCTTTGGGGGGCGGGGTCAGAAGCCAGAGCTGCAGATGCTCACTGTGTGCGAGGAGCTGCCGTTTGA  
CAACGTGGATGGTGGTGTGTGGAGGCAAGGCTTCGACATCTCCTACGACCCGCACGACTGGGATGCTGAA  
GACCTGCAGGTGTTTGTGGTGCCCCACTCTCACAATGACCCAGGCTGGATCAAGACCTTTGACAAGTACT  
ACACAGAGCAGACCCAACACATCCTCAATAGCATGGTGTCTAAGCTGCAGGAGGACCCCCGGCGCGCTT  
CCTCTGGGCAGAGGTCTCCTTCTTCGCCAAGTGGTGGGACAACATCAATGTCCAAAAGAGAGCGGCAGTC  
CGAAGGCTGGTGGGAAACGGGCAGCTGGAGATTGCGACAGGAGGCTGGGTGATGCCAGATGAGGCCAATT  
CCCCTACTTTTGCAATTGATTGACCAGCTCATCGAAGGACACCAGTGGCTGGAGAGAAATCTTGGTGCAAC  
CCCCCGCTCTGGCTGGGCAGTGGACCCCTTTGGATACAGCTCCACCATGCCTTACCTGCTGCGCCGTGCC  
AACCTCACCAGCATGCTGATTGAGAGAGTGCATATGCCATCAAGAAGCACTTTGCTGCCACCCACAGCC  
TAGAGTTTATGTGGAGGCAGACATGGGACTCGGACTCCAGCACAGACATCTTCTGTACATGATGCCCTT  
CTACAGCTATGACGTCCCCATACCTGTGGCCCAGATCCCAAGATCTGCTGCCAATTTGATTTCAAACGC  
CTGCCCTGGTGGGCGCATCAACTGCCCTTGGGAAGGTGCCACCCCGGGCCATCACAGAGGCCAACGTGGCAG  
AGAGGGCAGCCCTGCTTCTGGACCAATACCGGAAGAAGTCCAGCTGTTCCGAAGCAACGTCTCTCTGGT  
GCCTCTTGGAGATGACTTCCGATATGACAAGCCCCAGGAGTGGGATGCCAGTTCTTCAACTACCAACGG  
CTCTTTGACTTCTTCAACAGCAGGCCTAACCTCCATGTGCAGGCCAGTTTGGCACTCTTTCTGACTATT  
TTGATGCCCTGTACAAGAGGACAGGGGTGGAGCCAGGGGGCCCCGGCTCCAGGGTTTCTGTGCTGAGCGG  
GGATTTCTTCTCCTATGCGGACCGGGAGGATCATTACTGGACAGGCTATTACACTTCCCGGCCCTTCTAC  
AAGAGCTTAGACCGAGTCTTGAAGCCACCTGCGGGGGCAGAGGTCTGTACAGCCTGGCTGCAGCTC  
ACGCTCGCCGCTCTGGTCTGGCTGGCCGGTACCCACTGTCTGATTTCACCTCCTGACGGAAGCTCGGCG  
CACATTGGGGCTCTTCCAGCATCACGATGCCATCACTGGCACGGCCAAGGAGGCTGTGGTGGTGGACTAT  
GGGGTCAGGCTTCTGCGCTCCCTTGTCAACCTGAAGCAGGTCACTATTGATGCAGCCCACTATCTGGTGC  
TGGGGGACAAGGAGACTTACCCTTTGACCTGAGGCGCCCTTCTTCCAAGTGGATGACACTCGCTTAAG  
TCACGACGCCCCTCCAGAGCGCACGGTGTATCCAGTGGATTCTCGCCAGGTTTGTGGTCTATTCAAC  
CCACTGGAACAGGAGCGATTACAGCATGGTGTCCCTGTGTTCAACTCTCCCCGCGTGCCTGTCTTTTCGG  
AGGAGGGTCAGCCCTGGCCGTGCAGATCAGCGCACACTGGAGCTCTGCCACCGAGGCGGTCCCTGACGT  
CTACCAGGTGTCTGTGCCTGTCCGCCTGCCAGCCCTGGGCCGTGGGCGTGTGTCAGCTACAGCTGGGCCTG  
GATGGGCACCGCACGCTGCCCTCCTCTGTGCGCATCTACCTGCACGGCCGGCAGCTGTCCGTEAGCAGGC  
ACGAAGCGTTTCTCTCCGTGTCACTGACTCTGGCACCAGCGACTTCGCCCTCAGCAACCGCTACATGCA  
GGTCTGGTTCTCAGGCCCTTACTGGGCTCCTCAAGGGGTGAGGGCTGTGTTTTTTGGCAGAGCATCCGAAG  
GGTGGATGAGGAGCACGAGCAGCAGGTGGACATGCAGGTCTTGTCTATGGCACCCGTACGTCCAAAGAC  
AAGAGTGGAGCCTACCTCTTCTTCCGATGGCGAGGCAAGCCCTACGTCCCCAAGGAGCCCCCGTGTCT  
GCGTGTCACTGAAGGCCCTTTCTTCTCAGAGGTGGTTGCGTACTATGAGCACATTACAGGCGGTCCGG  
CTTTACAATCTGCCAGGGGTGGAGGGGCTGTCTCTGGACATATCATCCCTGGTGGACATCCGGGACTACG  
TCAACAAGGAGCTGGCCCTGCACATCCATACAGACATCGACAGCCAGGGTGCAGCCCCGACGGTATCTGA  
AGAAGCTCCCCCTCCAGGCCAACTTCTACCCCATGCCAGTCATGGCCTATATCCAGGACGCACAGAAGCG  
CCTCACGCTGCACACTGCCCAGGCCCTGGGTGTCTCTAGCCTCAAAGATGGCCAGCTGGAGGTGATCTTG  
GACCGGCGGCTGATGCAGGATGACAACCGGGGCTAGGCCAAGGGCTCAAGGACAACAAGAGAACCTGCA  
ACCGTTTCCGCCCTCTGCTAGAGCGGCGAACCCTGGGCAGTGAGGTCCAAGATAGCCACTCTACCAGCTA  
CCCATCCCTCCTCAGCCACCTGACCTCCATGTACCTGAACGCCCCGGCGCTCGCTCTGCCTGTAGCCAGG  
ATGCAGCTCCCAGGCCCTGGTCTGCGCTCATTTTCATCTCTGGCTTCTTCACTGCCCTGTGACTTCCACC  
TGCTCAACCTACGTACGCTCCAGGCTGAGGAGGCAACCTACCTCGGCGGAGACCGCACTCATCTTACA  
CCGCAAGGTTTGTGACTGCGGCCCTGGAGGCCAAGAACCTTGGGCTTCAACTGCACCACAAGCCAAGGCAAG  
GTAGCCCTGGGCGAGCCTTTTTCATGGCCTGGATGTGGTATTCTTTCAGCCAACCTCCTTGACGTTACTGT  
ACCTCTGGCCTCCCCGTCCAACAGCACTGACGTCTATTTGGAGCCCATGGAGATTGCTACCTTTTCGCCT  
CCGCTTGGGTTAGGGCTTCTTGTGGCCTGAAGAGAAAGTTTCAATTCACAGAGACTGCCTCTTAACATGAAG  
ATCATTTGGACAAGCCACACGGGTATCCCATCCCGATCTGCCTCCAGAACTGTGACACACTGGGCTCTGC  
CCTCATTTTCTGTTTTATTGCTGCTGCTGTGTTTTTCGGCGCAACCCACAAACCCAGTGATGGGTAAATAGG  
GCAGACGCCAGTGAGATCAGGGAGAGAAGGCCCTTGGTCAGAGTGGGCGAGTGCCAGGCTCTGCTTTGGGT  
TGTGAGTGGACACCCAAC'TGGGCACAGGCTCAGGCACCCATCCTTTTTTCCAAACAGGGATATAGAAGTGG

TGGAAGCAGACAGAAGAGGTAAGGGAGGCTAAGTGGGTAACAGCCCAGCATCAGGGTCACTGTGGCAACA  
GCAGGCTCTAGGGGAATCCTGTGGTTATGTAGAGACTCCATGTCCTGGTGTGATGAGCAGGATCAGAGTG  
ACTCTGGGAGGACAGGGGTGGGGACCCAGAGTTAGCAGTGGGGATGGAGCAGTAGAAGGAATCACTGTTT  
CTCCTAGGAGTCTGAAGGCCTCGCTGCTTTCTGTGATGGCTTTGCAGTAAGTGC CGCCTGGCCTGCATGC  
ATTGGCTAACAGGCTGCAGAATGGCAGGAAGGACTCGCTAGAGATTGTCATGGCCAGAGATCATAGGTCA  
CTTCAGGTAGCAAGACCCCTGGCAAACCTGGGCACCTGGCCTATGTACTGATTTGTGGGATGGTGGCAGGG  
GTGTGGGGTCCCTTACCCTGCCTGAATTCTCTTTGGCTTCTGTGCTCTGTATGCTGCTGTCCCAAGGGC  
TCTTTCTTATTATGGCAGGGAGTGGGGATTGGTCTACTTTCTTCTCTGGAAAGGAAAGCCTCCAAGAC  
TCCATGTGCTTGGGCAGCTTGAGAAGGCGTTACAGACCACGCC TAGCAGGCAGACCTTGAAGCCTCACCT  
TTAGTCTATCTGCAGAGGTATTAGTTCCTGGCACAGGGGACTAGGGGCATGTAGAGTATATGAGGAGGC  
AGTATGGCTGTGCAGGAGCCTTCATTTACAGTTCAATTAAATAGGGAAGAATTTATGATAGCTCTATAGAT  
GCTGAAAAGGTATTTCTGAAGATTTAAAATCCATCCCTTATTA AAACTCTTAGTAAATTAAGTCTGGAAA  
GAAACACCCTAATCTAGATAAAGGTCTGTTTCAGAAACCAACAGTGATGGCATTCTAAAGAGTCAGACGC  
CACAGGCATTCCCATTAAGTCAAGAACTAGCCAAGGGCAAGCTATTATTACAGCAGTGTCCCGGCACTAC  
TAACCCCTGCAACAAGCCAGATGAGGAACATAAGGAAGAATTATAATTGTCATTATTTGTAGACAATAAA  
ACTGCCTACCTGTAAAA

>gi|10835172|ref|NM\_000620.1| Homo sapiens nitric oxide synthase 1  
(neuronal) (NOS1), mRNA

AGAGCGGCTCTTTTAATGAGGGTTGCGACGTCTCCCTCCCCACACCCATAAACCAGTCGGGTTGGACGTC  
ACTGCTAATTCGTTTCAGTGATGATAGGATAAAGGAGGGACATTAAGAAATAAATTCCCCCTCACGACCC  
TCGCTGAGCTCACGGCTCAGTCCCTACATATTTATGCCGCGTTTCCAGCCGCTGGGTGAGGAGCTACTTA  
GCGCCGCGGCTCCTCCGAGGGGCGGCGGGCAGCGAGCAGCGGCCGAGCGGACGGGCTCATGATGCCTCA  
GATCTGATCCGCATCTAACAGGCTGGCAATGAAGATACCCAGAGAATAGTTCACATCTATCATGCGTCAC  
TTCTAGACACAGCCATCAGACGCATCTCCTCCCCCTTTCTGCCTGACCTTAGGACACGTCCCACCGCCTCT  
CTTGACGTCTGCCTGGTCAACCATCACTTCCTTAGAGAATAAGGAGAGAGGCGGATGCAGGAAATCATGC  
CACCAGCGGGCCACCAGCCATGAGTGGGTGACGCTGAGCTGACGTC AAAGACAGAGAGGGCTGAAGCCTT  
GTCAGCACCTGTCACCCCGGCTCCTGCTCTCCGTGTAGCCTGAAGCCTGGATCCTCCTGGTGAAATCATC  
TTGGCCTGATAGCATTGTGAGGTCTTCAGACAGGACCCCTCGGAAGCTAGTTACCATGGAGGATCACATG  
TTCGGTGTTCAGCAAATCCAGCCCAATGTCAATTTCTGTTCTGCTCTTCAAGCGCAAAGTTGGGGGCTGG  
GATTTCTGGTGAAGGAGCGGGTCAGTAAGCCGCCGTGATCATCTCTGACCTGATTCTGGGGGCGCCGC  
AGAGCAGAGAGCGCCTCATCCAGGCGGAGACATCATTTCTGCGGTCAACGGCCGGCCCTTGGTGACCTG  
AGCTATGACAGCGCCCTGGAGGTACTCAGAGGCATTGCTCTGAGACCCACGTGGTCTCATTTCTGAGGG  
GCCCTGAAGGTTTACCACGCACCTGGAGACCACCTTTACAGGTGATGGGACCCCCAAGACCATCCGGGT  
GACACAGCCCCCTGGGTCCCCCACC AAAGCCGTGGATCTGTCCACCAGCCACCGGCCGGCAAAGAACAG  
CCCCCTGGCAGTGGATGGGGCCTCGGGTCCCGGAATGGGCCTCAGCATGCCTACGATGATGGGCAGGAGG  
CTGGCTCACTCCCCCATGCCAACGGCTGGCCCCCAGGCCCCCAGGCCAGGACCCGCGAAGAAAGCAAC  
CAGAGTCAGCCTCCAAGGCAGAGGGGAGAACAATGAAC TGCTCAAGGAGATAGAGCCTGTGCTGAGCCTT  
CTCACCAGTGGGAGCAGAGGGGTCAAGGGAGGGGCACCTGCCAAGGCAGAGATGAAAGATATGGGAATCC  
AGGTGGACAGAGATTTGGACGGCAAGTCACACAAACCTCTGCCCCCTCGGCGTGGAGAACGACCGAGTCTT  
CAATGACCTATGGGGGAAGGGCAATGTGCCTGTCTGCTCTCAACAACCCATATTTCAGAGAAGGAGCAGCCC  
CCCACCTCAGGAAAACAGTCCCCCACAAGAATGGCAGCCCCCTCCAAGTGTCCACGCTTCTCAAGGTCA  
AGAACTGGGAGACTGAGGTGGTTCTCACTGACACCCTCCACCTTAAGAGCACATTGGAACGGGATGCAC  
TGAGTACATCTGCATGGGCTCCATCATGCATCCTTCTCAGCATGCAAGGAGGCCTGAAGACGTCCGCACA  
AAAGGACAGCTCTTCCCTCTCGCCAAAGAGTTTATTGATCAATACTATTTCATCAATTAAGAGATTTGGCT  
CCAAAGCCCACATGGAAGGCTGGAAGAGGTGAACAAAGAGATCGACACCACTAGCACTTACCAGCTCAA  
GGACACAGAGCTCATCTATGGGGCCAAGCACGCCTGGCGGAATGCCTCGCGCTGTGTGGGCAGGATCCAG  
TGGTCCAAGCTGCAGGTATTGATGCCCCGTGACTGCACCACGGCCCACGGGATGTTCAACTACATCTGTGA  
ACCATGTCAAGTATGCCACCAACAAAGGGAACCTCAGGTCTGCCATCACCATATTCCCCCAGAGGACAGA  
CGGCAAGCACGACTTCCGAGTCTGGAACCTCCAGCTCATCCGCTACGCTGGCTACAAGCAGCCTGACGGC  
TCCACCTGGGGGACCCAGCCAATGTGCAGTTTCAGAGATATGCATACAGCAGGGCTGGAAACCGCCTA  
GAGGCCGCTTCGATGTCTCGCGCTCCTGCTTTCAGGCCAACGGCAATGACCCTGAGCTCTTCCAGATTCC  
TCCAGAGCTGGTGTGGAAAGTTCCCATCAGGCACCCCAAGTTTGAGTGGTTCAAGGACCTGGGGCTGAAG  
TGGTACGGCCTCCCCGCGGTGTCCAACATGCTCCTAGAGATTGGCGGCCTGGAGTTACAGCGCCTGTCCCT  
TCAGTGGCTGGTACATGGGCACAGAGATTGGTGTCCGCGACTACTGTGACAACTCCCGCTACAATATCCT  
GGAGGAAGTGGCCAAGAAGATGAAC TTAGACATGAGGAAGACGTCCTCCCTGTGGAAGGACCAGGCGCTG  
GTGGAGATCAATATCGCGGTTCTCTATAGCTTCCAGAGTGACAAAGTGACCATTGTTGACCATCACTCCG

CCACCGAGTCCTTCATTAAGCACATGGAGAATGAGTACCGCTGCCGGGGGGGCTGCCCTGCCGACTGGGT  
 GTGGATCGTGCCCCCATGTCCGGAAGCATCACCCCTGTGTTCCACCAGGAGATGCTCAACTACCGGCTC  
 ACCCCCTCCTTCGAATACCAGCCTGATCCCTGGAACACGCATGTCTGGAAAGGCACCAACGGGACCCCCA  
 CAAAGCGGCGAGCCATCGGCTTCAAGAAGCTAGCAGAAGCTGTCAAGTTCTCGGCCAAGCTGATGGGGCA  
 GGCTATGGCCAAGAGGGTGAAAGCGACCATCCTCTATGCCACAGAGACAGGCAATCGCAAGCTTATGCC  
 AAGACCTTGTGTGAGATCTTCAAACACGCCTTTGATGCCAAGGTGATGTCCATGGAAGAATATGACATTG  
 TGCACCTGGAACATGAACTCTGGTCTTGTGGTCACCAGCACCTTTGGCAATGGAGATCCCCCTGAGAA  
 TGGGGAGAAATTCCGCTGTGCTTTGATGGAAATGAGGCACCCCAACTCTGTGCAGGAAGAAAGGAAGAGC  
 TACAAGGTCCGATTCAACAGCGTCTCCTCTACTCTGACTCCCAAAAATCATCAGGCGATGGGCCCCGACC  
 TCAGAGACAACCTTTGAGAGTGTGGACCCCTGGCCAATGTGAGGTTCTCAGTTTTTGGCCTCGGCTCAGC  
 AGCATACCCTCACTTTTGCGCCTTCGGACACGCTGTGGACACCTCCTGGAAGAAGCTGGGAGGGAGAGG  
 ATCCTGAAGATGAGGAAGGGGATGAGCTCTGTGGGCAGGAAGAGGCTTTTCAGGACCTGGGCGGAGAAGG  
 TCTTCAAGGCAGCCTGTGATGTCTTCTGTGTGGGAGATGATGTCAACATTGAAAAGGCCAACAATTCCCT  
 CATCAGCAATGATCGCAGCTGGAAGAGAAACAAGTTCCGCCTCACCTTTGTGGCCGAAGCTCCAGAAGCTC  
 ACACAAGGTCTATCCAATGTCCACAAAAGCGAGTCTCAGCTGCCCGGCTCCTTAGCCGTCAAAACCTCC  
 AGAGCCCTAAATCCAGTCGGTCAACTATCTTCGTGCGTCTCCACACCAACGGGAGCCAGGAGCTGCAGTA  
 CCAGCCTGGGGACCACTGGGTGTCTTCCCTGGCAACCACGAGGACCTCGTGAATGCCCTGATCGAGCGG  
 CTGGAGGACGCGCCGCTGTCAACCAGATGGTGAAAGTGGAAGTGTGGAGGAGCGGAACACGGCTTTAG  
 GTGTATCAGTAAGTGGACAGACGAGCTCCGCCTCCCGCCCTGCACCATCTTCAGGCCCTCAAGTACTA  
 CCTGGACATCACACGCCACCAACGCCCTCTGCAGCTGCAGCAGTTTGCCTCCC TAGCTACCAGCGAGAAG  
 GAGAAGCAGCGTCTGCTGGTCTCAGCAAGGGTTTGCAGGAGTACGAGGAATGGAATGGGGCAAGAACC  
 CCACCATCGTGGAGGTGCTGGAGGAGTTCCCATCTATCCAGATGCCGGCCACCCTGCTCCTGACCCAGCT  
 GTCCCTGCTGCAGCCCCGCTACTATTCCATCAGCTCCTCCCCAGACATGTACCCTGATGAAGTGCACCTC  
 ACTGTGGCCATCGTTTTCTACCGCACTCGAGATGGAGAAGGACCAATTACCCACGGCGTATGCTCCTCCT  
 GGCTCAACCGGATACAGGCTGACGAAGTGGTCCCCTGTTTTCTGTGAGAGGAGCACCCAGCTTCCACCTGCC  
 CCGGAACCCCCAAGTCCCCTGCACTCTCGTTGGACCAGGCACCGGCATTGCCCTTTCCGAAGCTTCTGG  
 CAACAGCGGCAATTTGATATCCAACACAAAGGAATGAACCCCTGCCCATGGTCTTGGTCTTCCGGGTGCC  
 GGCAATCCAAGATAGATCATATCTACAGGGAAGAGACCCTGCAGGCCAAGAACAAGGGGGTCTTCAGAGA  
 GCTGTACACGGCTTACTCCCGGGAGCCAGACAAACCAAGAAGTACGTGCAGGACATCTTGCAGGAGCAG  
 CTGGCGGAGTCTGTGTACCGAGCCCTGAAGGAGCAAGGGGGCCACATATACGTCTGTGGGGACGTACCA  
 TGGCTGCTGATGTCTCAAAGCCATCCAGCGCATATGACCCAGCAGGGGAAGCTCTCGGCAGCCAGCGC  
 CGGCGTATTTCATCAGCCGGATGAGGGATGACAAACCGATACCATGAGGATATTTTTGGAGTACCCCTGCGA  
 ACGTACGAAGTGAACACCGCCTTAGATCTGAGTCCATTGCCTTCATTGAAGAGAGCAAAAAAGACACCG  
 ATGAGGTTTTTCAGCTCCTAACTGGACCCCTCTTGCCAGCCGGCTGCAAGTTTTGTAAAGCGCGGACAGACA  
 CTGCTGAACCTTTCTCTGGGACCCCTGTGGCCCTCGCTCTGCCTCCTGTCTTGTCTGCTGTGCCCTGG  
 TTTCCCTCCTCGGGCTTCTCGCCCCCTCAGTGGTTTCTCGGCCCTCCTGGGTTTACTCCTTGAGTTTTCC  
 TGCTGCGATGCAATGCTTTTCTAATCTGCAGTGGCTCTTACAAAACCTGTCTCCCACTCCCTCTCTTGCC  
 GACAAGGGCAACTACGGGTGCATGAAACCACTGGAACATGGCCGTGCTGTGGGGGTTTTTTTCTCTGG  
 GGTTCCCCCTGGAAGGGCTGCAGGAAGTAGGCACAAGCTCTCTGAGCCAGTCCCTCAGCCACTGAAGTCCC  
 CCTTTCTCCTTTTTTATGATGACATTTTGGTTGTGCGTGCCTGTGTGTGTGTGTGTGTGTGTGTGTGT  
 GTGTGATGGGCCAGGTCTCTGTCCGTCTCTTCCCTGCACAAGTGTGTGATCTTAGATTGCCACTGCTT  
 TCATTGAAGACCCTCAATGCCAAGAAACGTGTCCCTGGCCCATATTAATCCCTCGTGTGTCCATAATTAG  
 GGTCCACGCCCATGTACCTGAAACATTTGGAAGCCCCATAATTGTTCTAGTTAGAAAGGGTTTCAGGGCAT  
 GGGGAGAGGAGTGGGAAATTGATTAAAGGGGCTGTCTCCCAATGAAAGAGGCATTCCCAAGATTTGCTGC  
 ATTTAGATTTTGATACCAGTGAGCAGAGCCCTCATGTGACATGAACCCATCCAATGGATTGTGCAAAATCC  
 CCTCCCCAAACCCACCCATACCAGCTAGAATCACTTGACTTTGCCACATCCATTGACTGACCCCTCCTC  
 CAGCAATAGCATCCAAGGGGCTGGAAGTTATGTTGTTCAAAGAAGCCTGGTGGCAATAAGGATCTTCCC  
 ACTTTGCCACTGGATGACTTTGGATGGGTCACTTGTCTCAGTTTTTCTTAGTCATAATGTATACGAAC  
 CTAAAGAATATGAATGGATTAAATGTTAAAGCTTTGGTGCCTGGAACAATATCAAGTAACAATATGATT  
 ATTATTTTTTTTATTTCCCCAAAGCGGGCTTGCTGCTTACCCCTTGGGGATGAAATAATGGAAGTGGTTA  
 AAGTGGATGAGGTGGAAGAGTTGCCATAATGAGTCCCACGTGGCTTCTTCGATAGGAGCCACAACCTT  
 GGGGTGGGAAGAAGCTTGTCCCTCAGGCTTGTGTCCTCTGCAGTTGATCTCCAAAGTTTTTAAACCTGTTA  
 AATTAATTTTGACAAATAAGTTACCCCTCAACTCAGATCAAAAATGGGCAGCCAAGTCTTCGGTAGGAATT  
 GGAGCCGGTGTAAATCCTCCCTAAGAGGCAACCTGTTGAATTTACTCTCTCAGAGTAAATGGTGGGAAGG  
 GATCCCTTTGTATACTTTTTTTAAATACTACAAATTAGTGTGAGGCAGTTCCAGAAAGAGACAAGAAATC  
 CTAGTGGCCTCCCAGACTGCAGGGTCCCCAAGGATGGAAGGGAATGTTCTGCTGGTTCTACCCTGTTTG  
 TTGTGTCTTGCTATACAGAAAAACCATTTCTTTTATATACTGTACGTGGGCATATCTTGTGTTTCACT

TTGGGTGTCTGCTAAAGAGGAAGTGCACCTGGCCCTCTTTGAAAGGGCTTTACAGTGGGGGCACCAAGACC  
 CCAAAGGCCCAGGCCAGGAGACTGTTAAAGTGAAAAGGCAATCTATGACTCACCTTGCTCTGCCATCCCT  
 GGCAGCCCCCACCAGGTGTCCTGTTCTGCCACATGGAGCTTGACTTCATGCCAGCTATAATCTCCCTGCT  
 CTTCTTTTAATCCCAATTTCCCTGCTCACTCTTCCACAGATATAAAGAACAACACTTAGCATCCCACA  
 CTCACCCCTTCTAATCCTGAAGGGAAGCCATTCTAAACTCCTTTCTGCAAACCCATTTCCAGCTCCTA  
 GTAGCTTTCTCCCAAAGGCTTTCTTTCCAATCCTTTATAGCTTTGGAGACGCCTCCCCAATTTCCCAAGG  
 GAAGGAAACTGTTGTGTCCAATCCCCATTAAAGACAAATTGATCAGTGCTTCCC

>gi|37540251|ref|XM\_291315.2| Homo sapiens hypothetical protein  
 FLJ23309 (FLJ23309), mRNA

GGCCGGGGCTGTCGCGGGTTGGGGCTGGCAGCTGAGGCTCGTGCCATGGAGTGGGGTTCTG  
 AGTCGGCTGTGTGAGGGCGCACGCGTCGGAGTAGAGCGTCGAGAGGGAGCGGCGGCCGCCACCGCC  
 GGAGAGGGAGGCGCGAGCGCAGGAGCCTCTGGTGGATGGGTGCAGCGCGGCGGGAGGACGCGGAAGAGG  
 AGCCCCGGGGGTAGCGGCGGCGCAGCAGGGGCGCGGGGACCGGGCTGTCTGAGGTGCGCGCCGCGCTGG  
 GGCTCGCGCTCTACCTGATCGCGCTGCGGACGCTGGTGCAGCTCTCGCTGCAGCAGCTCGTGCTACGCGG  
 GGCCGCTGGACACCGCGGGGAGTTCGACGCGCTCCAAGCCAGGGATTATCTTGAACACATAACCTCCATT  
 GGCCCCAGGACTACAGGAAGTCCAGAAAATGAAATTCTGACCGTGCACCTACCTTTTGAACAGATTAAAC  
 TGATTGAAGTGCAAAGCAACAGCCTTCATAAGATTTTCAGTAGATGTACAACGGCCACAGGCTCTTTTAG  
 CATTGATTTCTTGGGAGGTTTTACAAGCTATTATGACAACATCACCAATGTTGTGGTAAAGCTGGAACCC  
 AGAGATGGAGCCCAGCATGCTGTCTTGGCTAATTGTCATTTTGACTCAGTAGCAAACCTCACCAGGTGCCA  
 GTGATGATGCAGTTAGCTGCTCAGTGATGCTGGAAGTCCTTCGCGCTTGTCAACATCTTCAGAAGCCTT  
 GCATCATGCTGTCAATTTCTCTTTAATGGTGTGAGGAAAATGTCTTGCAAGCCAGTCATGGTTTCATT  
 ACTCAGCACCCCTGGGCTAGCTTGATTTCGTGCATTCAATTAACCTAGAGGCAGCAGGTGTAGGAGGGAAAG  
 AACTTGTATTCCAAACAGGTCTGAAAATCCTTGGTTGGTTCAAGCTTATGTTTCAGCAGCTAAACACCC  
 TTTTGCTTCTGTGGTGGCTCAGGAGGTTTTTCAGAGTGGAATCATTCCCTCAGATACTGACTTTTCGTATC  
 TACAGGGATTTTGGGAACATTCAGGAATAGACTTAGCTTTTATTGAGAATGGATACATTTATCACACCA  
 AGTATGACACAGCGGACAGAATTCCTAACAGATTCCATTACAGAGCAGGTGACAACATTTTAGCAGTTCT  
 TAAGCATCTAGCTACATCTGATATGCTGGCTGCTGCTTCTAAGTATCGACATGGAAACATGGTCTTCTTT  
 GATGTGCTGGGCCTGTTTGTCAATTGCCTACCCCTCTCGTATTGGCTCAATCATAAACTACATGGTGGTAA  
 TGGGTGTTGTTTTGTACCTGGGCAAAAAATTTTTGCAGCCCAACATAAGACTGGTAACATAAGAAGGA  
 CTTCTTGTGTGGACTTGGCATCACTTTGATCAGCTGGTTACTAGCCTTGTTACCGTTCTCATTATAGCA  
 GTGTTCACTCTCTCTTATTGGACAGTCTCTCTCATGTGTTAACCACTTCTATGTCTCCGTTTGTCTGTATG  
 GAACGTCAACTGTAGCCAAAATAATACTTATACATACTCTTGCGAAAAGATTTTATTACATGAATGCCAG  
 TGCCCAGTATCTGGGAGAAGTATTTTTTGACATTTGCTGTTTGTCCATTGCTGTTTTCTTGTACCCCTC  
 ACTTACCAAGGACTTTGCTCGGCGTTTATTAGTGCTGTCTGGGTAGCATTCCCATTGCTCACAAAGCTCT  
 GTGTGCATAAGGACTTCAAGCAGCATGGTGCCCAAGGAAAATTTATTGCTTTTTTACCTTTTGGGGATGTT  
 TATTCCTTATCTTTATGCATTGTACCTCATCTGGGCAGTATTGAGATGTTTACCCCTATCCTCGGGAGA  
 AGTGGTTCTGAAATCCCACCTGATGTTGTGCTGGCATCCATTTTGGCTGGCTGTACAATGATTCTCTCGT  
 CCTATTTTATTAACCTTCATCTACCTTGCCAAGAGCACAAAAAAACCATGCTAACTTTAACTTTGGTATG  
 TGCAATTACATTCTCTCTTGTTTGCAGTGGAACATTTTTTCCATATAGCTCCAATCCTGCTAATCCGAAG  
 CCAAAGAGAGTGTCTTTCAGCATATGACTAGAACATTCCATGACTTGGAAGGAAATGCAGTTAAACGGG  
 ACTCTGGAATATGGATCAATGGGTTTGATTATACTGGAATTTCTCACATAACCCCTCACATTCTGAGAT  
 CAATGATAGTATCCGAGCTCACTGTGAGGAGAATGCACCTCTTTGTGGTTTTCTTGGTATCTTCCAGTG  
 CACTTTCTGATCAGGAAAACTGGTATCTTCTGCCCCAGAAGTTTCTCCAAGAAATCTCTCTCATTTCC  
 GACTCATATCCAAAGAACAGACACCTTGGGATTCTATAAAATTGACTTTTGAAGCAACAGGACCAAGCCA  
 TATGTCCTTCTATGTTTCGAGCCCAAAAGGTCACACTTTCTCAGTGGTCTCTTGGCAATGGCACCCCA  
 GTCACAAGTAAAGGAGGAGACTACTTTGTCTTTTACTCCCATGGACTCCAGGCCTCTGCATGGCAGTTCT  
 GGATAGAAGTGACAGTTTTCAGAAGAACATCCTGAAGGAATGGTCACCGTGGCCATTGCTGCCCACTATCT  
 GTCTGGGGGAAGACAAGAGATCCCCCAACTGGATGCTCTGAAGGAAAAGTTCCAGATTGGACATTTCCC  
 TCTGCCCTGGGTGTGCACCTACGATCTCTTTGTATTTTAATCTTGTGGATGAGCTCTAAGTACATGCCAG  
 TGGACTATCCATGTGACATGGTTTCTCCCTATGTTTACGTGGATGTTTGTAAACGTAAGTCAATGAATTTTA  
 ATGATCATATGTTCAAAGAGCTTTCTGGGTTAACGCTTTTTCAGGGCCAAGCACTATAAGGGTTTAGCTGT  
 GGCGCAGTGATGCATGGCCTGTTGACACTTGAAAATGCCAGTCTTTTGGCACTTCAGCACATGTGGGTAC  
 TGCCACTACACACACGTCATTTTATATGACCTTAAGGACAAAGCCAACAATCCACTTCAATAGCTGCCCC  
 TTTAGGATCAAGAAAGATGTACACTGTACAGCATTGTTAATGAGACAAAAGTTGTTTCCAATTTAAGCC  
 CAAAACCATTTGTTGTATTAGTGGATGGTGGGTAAAATATCATTCACTGAGGTAATGATTCCCTTTGAG  
 AATATAACTCTGTGTAGGTCACTGGAAAGTGATTGCCATAGGGCTGGGAGAGAAGCATTGCACTCTTGAG

GCTGTAGCCTGTGTCAAGCTGTTTCTTCAGGCAGCCTCTCAAATGTGCTTTGTCTCTCTGTGCTGAGGCC  
 TGGACCCCTGTGCTGAGCTGGTGACTCACTGTCTCTGACAAGTGGACACACAGATGCACTGCTGTGCTGCTT  
 TCCTGAGGTGGTTTTCTATGCCTGTTTTCTCTGAAACATGTCTGTTACCCCTCTCCATCTTACCAAGTT  
 GAAAAGGGGAATATTTGGCCACATACCCCTCTGGTTTTCTGAGGTTCTTTTGGTTCAGAATATTGTTTGT  
 GCCAGTACATGACCTTAACCTTCTCTCAGAGCACTGAGCTGCCATCTGGGCTATTCTGGGGTAGAAGG  
 AAGGCTGGGAGTGGTGGGAATTTTATAAATATTTATCTCTTTTCTTTGTTTCATAGGAGTCTTGTGTTA  
 TACAAGGTTAGTCCTTCATGGTATAATCTTACTGATGCACTGGGCCATCTTTTGTTTTCCAGCCAGTT  
 GAATAGATTAGTTTTCTCAGTAACCTTACTATCCAGCAGACTGGCTTTCTTGAGACTTGAGGTTGTGGCT  
 TATACTGGAATGAGACCACGTACGTGTAGGTGGTTTCAGATCCTGCGTAATGGCAGCATGAGGACTTAAA  
 AGGTGGTTTTCTATTTGAAGATGGCTATGTAGCTTGTAAAGGTGATCACAGCAGTACCTCTCATGGCTTT  
 TTGGTTCCAGCAGTGGGGCATTTGGTGAGATCAATGGTAAACTGTGCAAGCTTTCTTTTTATCATTAGGA  
 AATGTGAAACGTTGGACAAATTTTGAGTTTTAACAAGGACAAAAAGTTGAAAGAAAAGGCACAGTTAACA  
 AAAAAGGGTGGCTAGATTTATCTTGGGTGATGGAGGAAATGAGAGAGGAATGCTCTTGAAAGGTGGTCTG  
 TGGATCTGTCTGAATAGAAAGAGCACAGTAAGTATGCATTGCCGGAGAAAACGTCCTTGAAGCTGCTTGT  
 CTCATGTGTATGATGTGCTTTTTAAATCATGCCCCCTCGTTGCCTGCCTAATCTGTGACTCCCTAAAACT  
 AACTGGGCCCATGTAGATGGGGCTGCAACCAGAGCTGAATAACATGTTAGGCTCACACATGCATCAGCAC  
 TGCACACTGGAATCATTGCTCTTCTGGACTTTGTAGAAATCAGTCTCAAGTGCTTCAAGAGTCTGGCTC  
 CTGCTACTTTTTATCTGTCTAGGTAGCACATAAGGTTTGCAGGGTTTATATTTTGTATAGAATCACAGTTGT  
 GGAGAAAAAGTAATAATTTCTCAATGAATTTAAAAATGGGCCATTTTCTATCCCCGTGGTTCATCTGA  
 TATAATTAGTGTTCCTGTGAATTCCTCCCCCTCTATGGGAAGGATGCCTTTACTCTTTATCAGTAATAAA  
 TTATGACTGTTTTCATATTGCCCTTAGGGTTATTTCCCTGTGTAAACCATTGTCTTTTGTTTTGGTTTTCT  
 TTAGCATTATGAAGCTTTGGTATTGTACAAGGTCAGTAGTAAGATGCTCACTAGTCTCAGGGCTTGTGTA  
 ATATTCTGGGAGGTCAATTTAAATGCCAGAAATGGTCAAGCAATTATACACAGTATTTATGACTCTGTTAA  
 GCATACCGTTTTGTCTGTCTACATTAGTAGATTCTGAGATTAAAAAAATTTTTAAAGAGTGATCATTTAAA  
 TAATTTCTAAAAGGGTCTTTTCAAGCTCTAACAAAGTCACCTAACAAATGCATTATTTTCTACAGAATTAG  
 ATGTTAGTAGTACAGTACTGCATATTCAGGGAAAAAGTGTGAGGAATTGATTTCAAATAGTTTCGTTCTT  
 GTGTTTGACCTAAGAATGATTGTTCGATGAAGTGTGTTTTTACAGTTTAGCATATATAAAACAAACATG  
 ATAGGATTCCTTAAGATGTTACCACCCAGGGGGCCACAAGCCAGCCTGCTGTCTCAGGAAGCTGTAGAAG  
 GAGTGTGTTGTCAATTTCTTGTCACTGGTTTGTCTGACTTACTGAGGATTAATTGTTGCCTTACAATGTTAC  
 TGAAATAAACTGTTTAATAT

>gi|21361381|ref|NM\_007171.2| Homo sapiens protein-O-  
 mannosyltransferase 1 (POMT1), mRNA

CCCGCAGGCTCGGTGAATCGAACGTTGAGCAGGGCGGTGGGTGGTGTGGAGTGCCGAGCGGCCTCACCCC  
 CAACCGTCGGGCCAGTCGGACGGTTCCGAGGCGTTGCCGGGAGCCGGGCGCGGCTCTGTGTGGACTCGGA  
 GAAACGCGGGGCGTCTGCCTGAGCCCGCTTTTCTACAAGATGTGGGGATTTTGAAGCGCCCTGTAGTGG  
 TGACGGCTGACATCAACTTGAGCCTTGTGGCCCTGACTGGGATGGGGTTACTGAGCCGGCTGTGGCGACT  
 CACCTACCCGCGGGCTGTGGTTTTTGTGACGAAGTATATTATGGGCAGTACATCTCTTTTACATGAAACAA  
 ATCTTCTTCTTGGATGACAGTGGGCCGCCATTTGGCCACATGGTGCTGGCCTTGGGAGGTTATTTAGGAG  
 GATTTCGATGGCAATTTTTTGTGGAACAGAATTGGAGCAGAATACAGTAGCAACGTGCCTGTGTGGTCCCT  
 GCGCCTGCTGCCAGCACTCGCGGGGGCCTTGTGGTCCCCATGGCCTACCAGATAGTGTGGAGCTCCAC  
 TTTTCTCATTTGTGCCGCCATGGGAGCTGCTCTGTTGATGCTTATCAAGAATGCTCTCATCACTCAGTCAA  
 GGCTAATGCTTTTGAATCAGTGTTAATATTTTTCAATCTATTGGCCGTGTTGTCTTACCTGAAGTTCTT  
 CAACTGCCAAAAGCACAGCCCTTTTTCTCTGAGCTGGTGGTTCTGGCTAACACTGACAGGGGTCTGCTTGT  
 TCCTGTGAGTGGGCATCAAGTACATGGGTGTGTTACGTACGTGCTCGTGCTGGGTGTTGCAGCTGTCC  
 ATGCCTGGCACCTGCTTGGAGACCAGACTTTGTCCAATGTAGGTGCTGATGTCCAGTGTGCATGAGGCC  
 GGCTGTATGGGGCAGATGCGGATGTACAGGGGTCTGTGTGTTCTGTCACTTGCTCGCCGAGCAGTG  
 GCTTTGCTGGTCATCCCGGTCGTCTGTACTTACTGTTCTTCTACGTCCACTTGATTCTAGTCTTCCGCT  
 CTGGGCCCCACGACCAATCATGTCCAGTGCCTTCCAGGCCAGCTTAGAGGGAGGACTAGCTCGGATCAC  
 CCAGGGTCAGCCACTGGAGGTGGCTTTGGGTCCCAAGGTACATCTGAGGAACGTCTTTGGGAAACCTGTG  
 CCTGCTGGCTTCATTCCCAACAGGACACCTACCCATGATATATGAGAACGGCCGAGGCAGCTCCCAAC  
 AGCAACAGGTGACCTGTTACCCCTTCAAAGACGTCAATAACTGGTGGATTGTAAAGGATCCCAGGAGGCA  
 CCAGCTGGTGGTGAGCAGCCCTCCGAGACCTGTGAGGCACGGGGACATGGTGCAGCTGGTCCACGGCATG  
 ACCACCCGCTCCCTGAACACGCATGATGTTGCAGCCCCCTGAGCCCCCATTCACAGGAGGTCTCCTGCT  
 ACATTGACTATAACATCTCCATGCCCCGCCAGAACCCTCTGGAGACTGGAAATTGTGAACAGAGGATCTGA  
 CACAGACGTCTGGAAGACCATCTCTCAGAGGTCCGCTTTGTGCACGTGAACACTTCCGCTGTCTTAAAG  
 CTGAGCGGGGCTCACCTCCCTGACTGGGGGTATCGGCAACTGGAGATCGTCGGGGAGAAGCTGTCCCGGG

GCTACCACGGGAGCACGGTGTGGAACTGAGGAGCACCGATACGGCGCGAGCCAGGAGCAGAGGGAGCG  
 GGAACGGGAGCTGCACCTGCAGAGGTGGACGTCAGCAGGAACCTCAGCTTCATGGCGAGATTCTCG  
 GAGCTGCAGTGGAGGATGCTGGCGCTGAGAAGTGATGACTCGGAACACAAGTACAGCTCCAGCCCACTGG  
 AGTGGGTACCCCTGGACACCAATATTGCCTACTGGCTGCACCCAGGACCAGCGCTCAGATCCACCTACT  
 TGGAAACATAGTGATCTGGGTTCGGGCAGCCTCGCTCTGGCCATCTACGCCCTGCTGTCTTGTGGTAC  
 CTGCTCCGACGGCGAAGAAATGTCCATGACCTCCCTCAGGATGCCTGGCTGCGCTGGGTGCTGGCTGGGG  
 CGCTGTGTGCCGTGGCTGGGCAGTGAACCTCCCGTTCTTCTGATGGAGAAGACACTCTTCTCTA  
 CCACTACCTGCCCGCACTCACCTTCCAAATCCTTCTGCTCCCTGTGGTCTTGCAGCACATCAGCGACCAC  
 CTGTGCAGGTCCCAGCTCCAGAGGAATTCTTTCAGCGCCCTGGTGGTGGCCTGGTACTCCTCCGCGTGCC  
 ACGTGTCCAAACAGCTGCGCCCACTCACTACGGGGACAAGTCACTCTCGCCACATGAACTCAAGGCCCT  
 TCGCTGGAAAGACAGCTGGGACATCTTGATCCGAAAACACTAGAACAAGAGTGTGGCAAAGAACACCCGT  
 GCTGGGGTCCGGACGAGGTTGAAGGGTCTTGGTCAATGTACGTAATGAGCAGGGTGGGCCACGCTGGGA  
 GGACACGGGTGGGCTGAGCAGGGCCTCTAGTGGAACACATTGGGGGTCTCATTGAAAAGCTCTCTGATG  
 AGCACCTCCTTTTGTGCAAAGTTAATTTTTTCTCGACAATAAAGATATTCCGTGTCTTACCCCTGAACT  
 AAGACACAGGGAGTATTTTCAAGAGCCAGCGTAGGAGTCATCGACAACGAAAAGCCGAGAACCAGGGCCA  
 GCAGTGGAAGCCTCAAGCAAGACCAGGGCCTGGTCTTGTCTAATTGCTGCAGGGTGGAGTTTGATCTGGC  
 AGACCCGATCCTCCTTTCATGAACACCCAGCAACCTGAGCAAGTCCCGGCCCTGCCCTCAGCGAGCCCGGC  
 AGGCGTCCCGGGACAGCTCAGTGTGGAGGGCCACCTGAACCACGAGCCAGGGCTGGGGCTTGATGTCA  
 TTGTCTATGACAGCGTCAAGACTGCCCTTGGCACCGTGTGTGTGGAAACCCTCCCTCTGAGACTCCAC  
 TGAGACGTGGCTGAGTGAATCTTCTCGTCAAGGTGTGTATCCATACAGCTCCATGCCTTT  
 GTCTTTTTTAAATGTAATTAATAAAGGAACCAACTGGCGTTTGTAAAAA

>gi|18860905|ref|NM\_002846.2| Homo sapiens protein tyrosine  
 phosphatase, receptor type, N (PTPRN), mRNA

GCGGCAAGCCCCGCTCAGCCCCCTTGGCAGGCTCCCGCCAGCGTCGCTGCGGCTCCGGCCCGGAGCGA  
 GCGCCCGGAGCTCGGAAAGATGCGGCGCCCGCGCGGCTGGGGGTCTCGGGGATCCGGGGGTCTCCGG  
 CTGCTCCTCTGCCTCTGCTGCTGAGCAGCCGCGGGGGGCTGCAGCGCGTTAGTGCCACGGCTGTC  
 TATTTGACCGCAGGCTCTGCTCTCACCTGGAAGTCTGTATTTCAGGATGGCTTGTGGGCAAGTGCAGGT  
 GGGAGTGGGGCAGGCCGGCCCCCTTTTGCAAGTCACTCCCAAGTTCTTCAACGCTTACAAGGTGTGCTC  
 CGACAACCTCATGTCCCAAGGATTGTCTGGCACGATGACCTACCCAGTATGTGATCTCTCAGGAGATGG  
 AGCGCATCCCCAGGCTTTCGCCCCCAGAGCCCCGTCCAAGGGACAGGTCTGGCTTGGCACCAGAGACC  
 TGGTCTGCTGGAGAGCTGCTTTTACAGGACATCCCCACTGGCTCCGCCCCCTGCTGCCCAGCATCGGCTT  
 CCACAACCACAGTGGGCAAAGGTGGAGCTGGGGCCAGCTCCTCTCTGTCCCCCTTGCAGGCTGAGCTGC  
 TCCCGCTCTCTTGGAGCACCTGCTGCTGCCCCCAGAGCTCCCCACCCTTCACTGAGTTACGAACCTGC  
 CTTGCTGCAGCCCTACCTGTTCCACAGTTTGGCTCCCGTGATGGCTCCAGGGTCTCAGAGGGCTCCCCA  
 GGGATGGTCAAGTGTGCGCCCCCTGCCCAAGGCTGAAGCCCCCTGCCCTCTTACAGCAGAACTGCCTCCAAGG  
 GCATATTTGGGGACCAACCTGGCCACTCCTACGGGGACCTTCCAGGGCTTCACTGCCCAGCTTTTTTCA  
 AGACTCTGGGTGCTCTATCTGGCCAGGAGTTGCCAGCACCCAGCAGGGCCAGGGTGC CAAGGTGCCA  
 GAGCAAGGGAGCAGCAGCCGGGCAGAGGACTCCCAGAGGGCTATGAGAAGGAAGGACTAGGGGATCGTG  
 GAGAGAAGCTGCTTCCCCAGCTGTGCAGCCAGATGCGGCTCTGCAGAGGCTGGCCGCTGTGCTGGCGGG  
 CTATGGGGTAGAGCTGCGTCAGCTGACCCCTGAGCAGCTCTCCACACTCTGACCCTGCTGCAGCTACTG  
 CCCAAGGGTGCAGGAAGAAATCCGGGAGGGGTGTAAATGTTGGAGCTGATATCAAGAAAACAATGGAGG  
 GGCCGGTGGAGGGCAGAGACACAGCAGAGCTTCCAGCCCGCACATCCCCATGCCTGGACACCCCACTGC  
 CAGCCCTACCTCCAGTGAAGTCCAGCAGGTGCCAAGCCCTGTCTCCTCTGAGCCTCCCAAAGCTGCCAGA  
 CCCCCCTGTGACACCTGTCTGTAGAGAAGAAAAGCCCACTGGGCCAGAGCCAGCCACGGTGGCAGGAC  
 AGCCCTCAGCCCGCCAGCAGCAGAGGAATATGGCTACATCGTCACTGATCAGAAGCCCTGAGCCTGGC  
 TGCAGGAGTGAAGCTGCTGGAGATCCTGGCTGAGCATGTGCACATGTCTCAGGCAGCTTCATCAACATC  
 AGTGTGGTGGGACCAAGCCCTCACCTTCCGCATCCGGCACAAATGAGCAGAACCTGTCTTTGGCTGATGTGA  
 CCCAACAAGCAGGGCTGGTGAAGTCTGAAGTCAAGAGCAGAGAGGGCTCCAAATCTTGACAGCAGGAGT  
 GGGACAGAGGGAGGAGGAGCTGCAGTCACTTCCCCAACTGCGCACAGCACCTACCCATGCGCTCAGTG  
 CTGCTCACTCTGGTGGCCCTGGCAGGTGTGGCTGGGCTGCTGGTGGCTCTGGCTGTGGCTCTGTGTGTGC  
 GGCAGCATGCGCGGCAGCAAGACAAGGAGCGCTGGCAGCCCTGGGGCTGAGGGGGCCCATGGTGACAC  
 TACCTTTGAGTACCAGGACCTGTGCCGCCAGCACATGGCCACGAAGTCTTGTTCACCGGGCAGAGGGT  
 CCACCGGAGCCTTACCGGTGAGCAGTGTGTCTCCAGTTCAGCGACGACAGCCAGGCCAGCCCACTGCT  
 CCCACAGCAGCACCCCGTCTTGGTGCAGGAGCCGGCCCAAGCCAACATGGACATCTCCACGGGACACAT  
 GATTCTGGCATAACATGGAGGATCACCTGCGGAACCGGGACCGCTTGCCAAGGAGTGGCAGGCCCTCTGT  
 GCCTACCAAGCAGAGCCAAACACCTGTGCCACCGCGCAGGGGGAGGGCAACATCAAAAAGAACCGGCATC

CTGACTTCCTGCCCTATGACCATGCCCCGATAACTGAAGGTGGAGAGCAGCCCTTCTCGGAGCGATTA  
CATCAACGCCAGCCCCATTATTGAGCATGACCCTCGGATGCCAGCCTACATAGCCACGCAGGGCCCGCTG  
TCCCATACCATCGCAGACTTCTGGCAGATGGTGTGGGAGAGCGGCTGCACCGTCATCGTCATGCTGACCC  
CGCTGGTGGAGGATGGTGTCAAGCAGTGTGACCGCTACTGGCCAGATGAGGGTGCCTCCCTCTACCACGT  
ATATGAGGTGAACCTGGTGTGCGAGCACATCTGGTGCAGGACTTCTGGTGCAGGCTTCTACCTGAAG  
AACGTGCAGACCCAGGAGACGCGCACGCTCACGCAGTTCCACTTCCTCAGCTGGCCGGCAGAGGGCACAC  
CGGCCTCCACGCGGCCCTGCTGGACTTCCGCAGGAAGGTGAACAAGTGCTACCGGGCCGCTCCTGCCC  
CATCATCGTGCAGTGCAGTGATGGTGCAGGAGGACCGGCACCTACATCCTCATCGACATGGTCCTGAAC  
CGCATGGCAAAAGGAGTGAAGGAGATTGACATCGCTGCCACCCTGGAGCATGTCCGTGACCAGCGGCTG  
GCCTTGTCCGCTCTAAGGACCACTTTGAATTTGCCCTGACAGCCGTGGCGGAGGAAGTGAATGCCATCCT  
CAAGGCCCTGCGCCAGTGAGACCCTGGGGCCCCCTGGCGGGCAGCCAGCCTCTGTCCCTCTTTGCCTGT  
GTGAGCATCTCTGTGTACCCACTCCTCACTGCCCCACCAGCCACCTCTTGGGCATGCTCAGCCCTTCCTA  
GAAGAGTCAGGAAGGGAAAGCCAGAAGGGGCACGCCTGCCAGCCTCGCATGCCAGAGCCTGGGGCATCC  
CAGAGCCCAGGGCATCCCATGGGGGTGCTGCAGCCAGGAGGAGAGGAAAGGACATGGGTAGCAATTCTAC  
CCAGAGCCTTCTCCTGCCTACATTCCCTGGCCTGGCTCTCCTGTAGCTCTCCTGGGGTTCTGGGAGTTCC  
CTGAACATCTGTGTGTGTCCCCCTATGCTCCAGTATGGAAGAATGGGGTGGAGGGTCGCCACACCCGGCT  
CCCCCTGCTTCTCAGCCCCGGGCTGCCTCTGACTCACACTTGGGCGCTCTGCCCTCCCTGGCCTCACGC  
CCAGCCTCCTCCCACCACCCTCCCACCATGCGCTGCTCAACCTCTCTCCTTCTGGCGCAAGAGAACATTT  
CTAGAAAAAACTACTTTTGTACCAGTGTGAATAAAGTTAGTGTGTTGTCTGTGCAGCTGCAAAAAAAA  
AAAAAAA

>gi|19743910|ref|NM\_002847.2| Homo sapiens protein tyrosine  
phosphatase, receptor type, N polypeptide 2 (PTPRN2), transcript  
variant 1, mRNA

GCCTCCCGCCGCTCCCGCGCGGCCATGGACTGAGCGCCGCCGCGCCAGGCCGCGGGGATGGGGCCGCCG  
TCCCGCTGCTGCTGCTGCTACTGCTGCTGCTGCCGCCACGCGTCTGCTGCCGCCCTTCTGCTCCGTC  
CCGCGGCCGCGCAGCTCCCGGGCGTCTGGGCTGCCTGCTCGAGGAGGGCCTCTGCGGAGCGTCCGAGGCC  
TGTGTGAACGATGGAGTGTGTTGGAAGGTGCCAGAAGGTTCCGGCAATGGACTTTTACCCTACGAGGTGT  
CGCCCGTGGCCCTGCAAGCGCTGCGCGTGGCGTTGCAGAAGCTTTCGGCACAGGTTTACGCTGGCAGGA  
TGACTATACTCAGTATGTGATGGACAGGAACCTTGCAGACCTCCGAAACCTACCTGAGGCGCTCTGAA  
GCATCCAGCCCAGCCAGGCCCTCAAAACACAGCGTTGGCAGCGAGAGGAGGTACAGTCGGGAGGGCGGTG  
CTGCCCTGGCCAACGCCCTCCGACGCCACCTGCCCTTCTTGGAGGCCCTGTCCAGGCCCCAGCCTCAGA  
CGTGCTCGCCAGGACCCATACGGCGCAGGACAGACCCCCGCTGAGGGTGATGACCGCTTCTCCGAGAGC  
ATCCTGACCTATGTGGCCACACGTCTGCGCTGACCTACCTTCCCGGGCCCCGACCCAGCTCCGCGAGG  
ACCTCCTGCCGCGGACCTCGGCCAGCTCCAGCCAGATGAGCTCAGCCCTAAGGTGGACAGTGGTGTGGA  
CAGACACCATCTGATGGCGGCCCTCAGTGCCTATGCTGCCAGAGGCCCCAGCTCCCCCGGGGAGGGC  
AGCCTGGAGCCACAGTACCTTCTGCGTGCACCTCAAGAATGCCAGGCCTTTGCTGGCACCAGCCGCCC  
CCCAGAAGTGGCCTTACCTCTGGGAGATTCCGAAGACCCCTCCAGCACAGGCATGGAGCACGGATTCA  
TACCTCCTGAAGGACCTGCAGAGGCAGCCGGCTGAGGTGAGGGGCTGAGTGGCCTGGAGCTGGACGGC  
ATGGCTGAGCTGATGGCTGGCCTGATGCAAGGCGTGACCATGGAGTAGCTCGAGGCAGCCCTGGGAGAG  
CGGCCCTGGGAGAGTCTGGAGAACAGGCGGATGGCCCCAAGGCCACCTCCGTGGAGACAGCTTTCCAGA  
TGACGGAGTGCAGGACGACGATGATAGACTTTACCAAGAGGTCCATCGTCTGAGTGCCACACTCGGGGGC  
CTCCTGCAGGACCACGGGTCTCGACTCTTACCTGGAGCCCTCCCTTTGCAAGGCCCTCGACATGGAGA  
GGAAGAAGTCCGAGCACCTGAGTCTTCCCTGTCTTCAAGAAGAGGAGACTGCCGGAGTGGAGAACGTCAA  
GAGCCAGACGTATTCCAAAGATCTGCTGGGGCAGCAGCCGATTCGGAGCCCGGGCCGCTGCGTTTGGG  
GAGCTCCAAAACAGATGCCTGGGCCCTCGAAGGAGGAGCAGAGCCTTCCAGCGGGTGCTCAGGAGGCC  
TCAGCGACGGCTGCAATTGGAGGTCCAGCCTTCCGAGGAAGAGGCGCGGGCTACATCGTGACAGACAG  
AGACCCCTGCGCCCCGAGGAAGGAAGCGGCTGGTGAGGACGTCGCGCGCTCTCTGAGGTGCCAGC  
AGTGCCCTTCGCTGACGTGGAGGTTCTCGGACCAGCAGTGACCTTCAAAGTGAGCGCCAATGTCCAAAACG  
TGACCATTGAGGATGTGGAGAAGGCCACAGTTGACAACAAAGACAACTGGAGGAAACCTCTGGAGTGA  
AATTCTTCAAACCGGAGTCGGGTGAAAAGCAAACCTCAAGTTCTTGCCTCCTCAGGCGGAGCAAGAAGAC  
TCCACCAAGTTCATCGCGCTCACCTGGTCTCCCTCGCTGCATCCTGGGCGTCTCCTGGCCTCTGGCC  
TCATCTACTGCCTCCGCCATAGCTCTCAGCACAGGCTGAAGGAGAAGCTCTCGGGACTAGGGGGCGACCC  
AGGTGCAGATGCCACTGCCGCTTACCAGGAGCTGTGCCGCCAGCGTATGGCCACGCGGCCACCAGACCGA  
CCTGAGGGCCCGCACACGTCACGCATCAGCAGCGTCTCATCCAGTTTACGCGACGGGCGGATCCCCAGCC  
CCTCCGCACGCAGCAGCGCTCATCCTGGTCCGAGGAGCCTGTGCAGTCCAACATGGACATCTCCACCGG  
CCACATGATCCTGTCTTACATGGAGGACCACCTGAAGAACAAGAACC GGCTGGAGAAGGAGTGGGAAGCG



CTGTGCGCCTACCAGGCGGAGCCCAACAGCTCGTTCTGTGGCCCAGAGGGAGGAGAACGTGCCCCAAGAACC  
GCTCCCTGGCTGTGCTGACCTATGACCACTCCCGGGTCTGTGCTGAAGGCGGAGAACAGCCACAGCCACTC  
AGACTACATCAACGCTAGCCCCATCATGGATCACGACCCGAGGAACCCCGCGTACATCGCCACCCAGGGA  
CCGCTGCCCCGCCACCGTGGCTGACTTTTGGCAGATGGTGTGGGAGAGCGGCTGCGTGGTGATCGTCATGC  
TGACACCCCTCGCGGAGAACGGCGTCCGGCAGTGCTACCACTACTGGCCGGATGAAGGCTCCAATCTCTA  
CCACATCTATGAGGTGAACCTGGTCTCCGAGCACATCTGGTGTGAGGACTTCCTGGTGAGGAGCTTCTAT  
CTGAAGAACCTGCAGACCAACGAGACGCGCACCGTGACGCAGTTCCTACTTCCTGAGTTGGTATGACCGAG  
GAGTCCCTTCCTCCTCAAGGTCCCTCCTGGACTTCCGCAGAAAAGTAAACAAGTGCTACAGGGGCCGTTT  
TTGTCCAATAATTGTTTCATTGCAGTGACGGTGCAGGCCGGAGCGGCACCTACGTCTGTATCGACATGGTT  
CTCAACAAGATGGCCAAAGGTGCTAAAGAGATTGATATCGCAGCGACCTGGAGCACTTGAGGGACCAGA  
GACCCGGCATGGTCCAGACGAAGGAGCAGTTTGAGTTCGCGCTGACAGCCGTGGCTGAGGAGGTGAACGC  
CATCCTCAAGGCCCTTCCCCAGTGAGCGGCAGCCTCAGGGGCCCTCAGGGGAGCCCCACCCACGGATGT  
TGTCAGGAATCATGATCTGACTTTAATTGTGTGCTTCTATTATAACTGCATAGTAATAGGGCCCTTAGC  
TCTCCCGTAGTCAGCGCAGTTTAGCAGTTAAAGTGTATTTTTGTTTAATCAAACAATAATAAGAGAGA  
TTTGTGGAATAATCCAGTTACGGGTGGAGGGGAATCGGTTTCATCAATTTTCACTTGCTTAAAAAATAC  
TTTTTCTTAAAGCACCCGTTACCTTCTTGGTTGAAGTTGTGTTAACAATGCAGTAGCCAGCACGTTCTGA  
GGCGGTTTCCAGGAAGAGTGTGCTTGTCTATCTGCCACTTTCGGGAGGGTGGATCCACTGTGCAGGAGTGG  
CCGGGGAAGCTGGCAGCACTCAGTGAGGCCGCCCGGCACACAAGGCACGTTTGGCATTCTCTTTGAGAG  
AGTTTATCATTTGGGAGAAGCCGCGGGGACAGAACTGAACGTCTTCAGCTTCGGGGCAAGTGAGACAATC  
ACAGCTCTCGCTGCGTCTCCATCAACACTGCGCCGGGTACCATGGACGGCCCCGTCAGCCACACCTGTC  
AGCCCAAGCAGAGTGATTACAGGGCTCCCCGGGGGAGACACCTGTGCACCCCATGAGTAGTGCCCACTT  
GAGGCTGGCACTCCCCTGACCTCACCTTTGCAAAGTTACAGATGCACCCCAACATTGAGATGTGTTTTTA  
ATGTTAAATATTGATTTCTACGTTATGAAAACAGATGCCCCCGTGAATGCTTACCTGTGAGATAACCAC  
AACCAGGAAGAACAATCTGGGCATTGAGCAAGCTATGAGGGTCCCCGGGAGCACACGAACCTGCCAGG  
CCCCCGCTGGCTCCTCCAGGCACGTCCCGGACCTGTGGGGCCCCAGAGAGGGGACATTTCCCTCCTGGGA  
GAGAAGGAGATCAGGGCAACTCGGAGAGGGCTGCGAGCATTTCCCTCCCGGGAGAGGAGATCAGGGCGAC  
CTGCACGCACTGCGTAGAGCCTGGAAGGGAAGTGAGAAACCAGCCGACCGGCCCTGCCCTCTTCCCGGG  
ATCACTTAATGAACCACGTGTTTTGACATCATGTAAACCTAAGCACGTAGAGATGATTCCGATTTGACAA  
AATAACATTTGAGTATCCGATTCCGCATCACCCCTACCCAGAAATAGGACAATTCACTTTCATTGACCA  
GGATGATACATGGAAGGCGGCGCAGAGGCAGCTGTGTGGGCTGCAGATTTCTGTGTGGGTTCCAGCGT  
AGAAAACGCACCTCCATCCCGCCCTTCCACAGCATTTCTCCATCTTAGATAGATGGTACTTCCAAAGG  
CCCTACCAGAGGGAACACGGCCTACTGAGCGGACAGAATGATGCCAAAATATTGCTTATGTCTCTACATG  
GTATTGTAATGAATATCTGCTTTAATATAGCTATCATTTCTTTTCCAAAATTACTTCTCTCTATCTGGAA  
TTTAATTAATCGAAATGAATTTATCTGAATATAGGAAGCATATGCCTACTTGTAATTTCTAACTCCTTAT  
GTTTGAAGAGAAACCTCCGGTGTGAGATATACAAATATATTTAATTGTGTATATTAACTTCTGATTCA  
AAAAAA

>gi|4507334|ref|NM\_003895.1| Homo sapiens synaptojanin 1 (SYNJ1), mRNA  
GAGGAGGAGCCGGAGGAAGAGGGCTGCCTCCGAAGAAAGGAGAATGGCGTTTCAGTAAAGGATTCGGGATC  
TATCACAATTTGGATCCCCACCTTTTACGCTCATAGTGGAACCTAGGCATAAGGAAGAATGTCTCATGT  
TCGAGTCTGGGGCTGTGCTGTGCTCTCATCTGCAGAAAAAGAGGCAATCAAGGGTACATACTCCAAAGT  
ACTGGATGCATATGGACTCTTAGGTGTTCTGCGGTTAAATCTTGGTGATACTATGTTACATTATCTGGTC  
CTAGTCACTGGATGTATGTCTGTTGGAAAAATTCAGAATCTGAAGTTTTCCGAGTTACTTCCACTGAGT  
TTATATCACTGCGAATCGATTCTTCAGATGAGGATCGCATTTTCAAGAGTGCGGAAAGTTTTGAATTCAGG  
AACTTTTATTTTGCATGGTCTGCATCTGGCATCAGTTTAGATTTGAGTCTTAATGCGCATCGTAGCATG  
CAAGAACAGACAACTGATAATAGATTTTTCTGGAATCAGTCTTTCGATTTGCATCTCAAACACTATGGCG  
TGAATTGTGATGACTGGTTATTACGTCTTATGTGTGGAGGAGTAGAAATCAGAACAAATTTATGCTGCTCA  
TAAACAGGCGAAGGCTTGCCCTCATTTCAAGATTTAAGCTGTGAACGAGCTGGGACCAGGTTTAAATGTCCGG  
GGAACAAATGATGATGGTCATGTTGCCAATTTTGTAGAAACAGAACAGGTTGTGTACTTAGATGACTCAG  
TTTCTTCCCTTACATACAAATCCGAGGATCTGTTCCATTGTTCTGGGAGCAACCAGGGTTGCAAGTGGGATC  
TCATCGTGTCCGTATGTCAAGGGGATTTGAAGCCAATGCACCTGCTTTTGCAGGCATTTTGAACACTT  
AAGAACTTATATGGTAGACAAATAATAGTAAATTTGCTTGGATCTAAGGAAGGTGAACATATGCTAAGTA  
AAGCTTTCCAGAGTCATTTGAAAGCTTCTGAACATGCTGCTGATATCCAGATGGTGAATTTTGAATATCA  
TCAATGGTTAAGGGAGGAAAGGCAGAAAAATTACATAGTGTCTTAAACCTCAAGTCCAGAAGTTTCTA  
GATTATGGATTTTTTTTATTTCAATGGAAGTGAAGTTCAAAGATGCCAGAGTGGTACAGTTTCAACAACT  
GCTTGGATTGTCTTGATAGAACAAATAGTGTGCAGGCATTTCTTGGCTTAGAGATGCTAGCTAAACAGTT  
GGAAGCTCTTGGTTTAGCTGAAAAGCCTCAGTTGGTGACTCGCTTTTCAAGAAGTTTTTCGGTCAATGTGG



TCCGTGAATGGTGATTCAATCAGTAAGATATATGCAGGAACCTGGAGCTCTTGAAGGGAAAGCGAAGTTAA  
AAGATGGTGCTCGCTCTGTTACCCGAACAATTCAGAATAACTTCTTTGACAGCTCCAAGCAAGAGGCCAT  
TGATGTTTTGCTACTGGGAAATACTCTGAATAGTGATTTAGCTGACAAAGCTCGAGCACTTTTAACTACT  
GGAAGTTTGGTGTCTTCTGAGCAGACATTACAGTCAGCATCTTCTAAAGTACTAAAGAGCATGTGTGAGA  
ATTTCTACAAATATTCAAAGCCTAAGAAAATTTCAGTATGTGTGCGAACCTGGAATGTGAATGGTGGGAA  
GCAATTTTCGAGCATAGCTTTTAAAGAATCAGACACTCACTGACTGGCTTCTTGATGCACCCAAGTTAGCT  
GGCATCCAGGAGTTTCAAGATAAAAGAAGTAAGCCAACCTGATATATTTGCAATTGGTTTTGAAGAAATGG  
TAGAATTGAATGCTGGAAACATTGTGAGTGCAAGCACAAACAAATCAGAAGCTCTGGGCTGTAGAACTTCA  
GAAGACAATCTCCAGAGACAACAAGTATGTGCTGCTGGCTTCTGAACAGTTGGTGGGCGCTGTTTTGTTT  
GTTTTTATCAGACCAGCATGCTCCTTTTATCAGGGATGTTGCAGTTGATACTGTGAAGACTGGAATGG  
GAGGTGCAACTGGAAATAAGGGAGCAGTTGCAATCCGAATGCTCTTCCATACAACCAGCCTTTGCTTCGT  
CTGTAGCCACTTTGCTGCAGGGCAGTCACAAGTCAAAGAAAAGAAATGAAGATTTTATAGAAATAGCACGA  
AAATTGAGTTTTTCTATGGGAAGGATGCTATTTTCCCATGACTATGTATTTTGGTGTGGTGATTTCAACT  
ATCGAATCGATCTCCCTAACGAAGAAGTTAAAGAGCTCATAAGACAGCAAAATTGGGATTCTCTTATAGC  
AGGAGATCAACTTATCAATCAGAAAAATGCTGGACAGGTTTTTATAGAGGATTTTATAGAAGGAAAGGTAACC  
TTTGCTCCGACATATAAGTATGACTTGTTTTCTGACGACTATGACACCAGTGAAAAGTGCCGCACCCCTG  
CCTGGACAGACCGTGTCTTTGGAGAAGGAGGAAATGGCCTTTTGATAGATCAGCTGAAGATCTAGATCT  
TCTAAATGCTAGTTTTCAAGATGAAAGCAAAATTCTGTACACGTGGACTCCAGGCACCTTTGCTGCACTAT  
GGAAGAGCTGAGCTGAAGACTTCTGACCACAGGCCGTGTCGTTGCCCTGATTGATATAGATATATTTGAAG  
TTGAAGCTGAAGAGAGGCAAAACATTTATAAGAAGTAATTGCAGTTCAGGGTCCACCAGATGGTACAGT  
ATTGGTCTCAATCAAAAGTTCTTTACCAGAAAATAATTTTTTTGATGATGCCTTGATTGATGAGCTTCTG  
CAGCAGTTTGCAAGTTTTGGTGAAGTTATACTTATAAGATTTGTAGAAGATAAAATGTGGGTACATTTT  
TGGAGGGAAGCTCTGCCTTGAATGTTCTGAGCCTAAATGGTAAAGAGTTATTGAATCGGACTATAACTAT  
TGCTTTAAAAAGTCCAGACTGGATCAAAAATTTGGAAGAAGAAATGAGTTTAGAGAAAATTAGCATTGCA  
TTGCCATCATCAACAAGCTCTACCCTGCTTGGTGAAGATGCAGAGGTTGCAGCAGATTTTGATATGGAAG  
GTGATGTTGATGACTATAGTGCTGAAGTGGAGGAACCTTCTCCTCAGCATCTCCAGCCATCTTCAAGTTC  
CGGCCTTGGTACTTCCCCCAGCTCTTACCCCGAAGTCTCCTGCCAGTCACCTACAATATCAGAGGGT  
CCTGTACCTTCCCTTCCCATCAGACCAAGCCGAGCACCCTCAAGAACTCTTGGGCTCTCCAGTGCACAGA  
GTTCTCCTATTGACGCGCAGCCAGCAACGCCGCTGCCGAGAAAGACCCGCCCCAGCCCTTGGAGCCCAA  
GCGGCCGCCGCCGCCGCCGCCGCTGCCCTCCACACGCCCGGCTCCCCACAGAGACCTCCTCCGCCCT  
TCAGGGGCTAGGATCTTGCACCCACTAGAAAGGAATTTGGAGGTATTGGAGCCCTCCAGTCTGGGG  
TAGCTAGGAGAGAGATGGAAGCACCCAAAAGCCCTGGAACAACAAGGAAAGATAATATAGGACGCAGTCA  
GCCTTACCTCAAGCAGGACTTGCAGGCCCAGGACCTGCTGGATACAGTACAGCCAGACCGACGATTCTCT  
CCTCGTGCTGGAGTTATCAGTGCCCCACAGAGCCACGCGCGGGCATCTGCTGGAAGACTGACTCCTGAAA  
GCCAAAGCAAAACATCAGAAACGTCGAAAGGTTCAACTTTCTTCTGAACCACTGAAGCCTCAGGCTGC  
TTTTCTCCTCCGAGTCTTCTTTGCCCCGCTGCTCAAAGGTTGCAAGAGCCTCTTGCTCCTGTGGCAGCA  
CCTATGCCTCAGTCTGGCCCCCAGCCAAATTTGGAACCCACCACAACCACCTCGAAGCAGGTCAT  
CCCATAGCTTGCTTTCAGAAGCTTCTTCAACCCGCAAGTAAAAACAAATGGAATCTCTGATGGCAAAAG  
AGAATCACCATTAAAGATTGACCCATTTGAAGATCTGTCATTTAATCTGCTTGCTGTATCAAAGGCTCAG  
CTATCTGTTCAAACGTCACCTGTTCCACCCAGACCCAAAGAGGTTGATTGAGTTGCCTTCTGCAACGC  
AAAGTAATGTTAATACTTTGAGTTCTGTAAGTTGCATGCCAACAATGCCTCCAATTCCAGCTCGGAGTCA  
ATCCCAGGAAAATATGCGAAGTTCTCCAAACCCATTTATTACTGGCTTGACCAGGACAAATCCTTTTCACT  
GACAGGACTGCTGCTCCTGGAAACCCATTTAGAGCCAAGTCTGAAGAATCAGAGGCAACTTCATGGTTCT  
CCAAAGAAGAGCCCGTTACTATCAGTCCTTTCCCTTCTTGCAGCCTCTTGGTCATAACAAAAGCAGGGC  
TTCATCTTCACTTGATGGCTTTAAGGACAGTTTTGATCTACAGGGCCAGTCTACATTAATAAATTAGCAAC  
CCGAAAGGATGGTAACCTTCGAGGAAGAAGAGGATTTTGGTGTGAAAGGGAAGTCAAAGTCAGCTTGTT  
CAGACTTACTGGGTAATCAGCCAAGTTCAATTTCTGGCTCCAACCTGACATTGAATGATGACTGGAATAA  
AGGGACAAATGTCTCCTTCTGTGTGTTGCCGTCAAGAAGACCTCCTCCACCTCCTGTCCCTCTGCTCCCG  
CCCGGCACCAGCCCTCCAGTAGATCCTTTCACGACCTTGGCCTCTAAGGCTTCACCCACACTGGACTTTA  
CAGAAAGATAACGCCAGCCCG

>gi|26190607|ref|NM\_003898.1| Homo sapiens synaptojanin 2 (SYNJ2), mRNA  
GGCACGAGGCGGGGAGCTGTGCGGGCAGCGGCCCTCGGGAGGACGTGGCCCCGCCCCCGCCGCGAGT  
GGGCCCCGACCCTCATGGCCCTGAGCAAAGGGCTGCGGCTGCTGGGGCGCCTGGGGGCCGAGGGGACTGT  
AGCGTGCTGCTGGAGGCGCGGGCCGCGACGACTGCCTGCTGTTTCGAGGCCGGCACGGTGGCCACGCTGG  
CTCCAGAAGAAAAGGAAGTCATTAAAGGACAGTATGGCAAGCTCACGGACGCGTACGGCTGCCTGGGGGA  
GCTGAGGCTGAAATCTGGTGGCACGTCTCTGAGCTTCTGGTGTGGTGACAGGCTGCACATCTGTGGGC

AGAATTCCAGATGCTGAAATCTACAAAATCACTGCCACTGACTTTTACCCTCTTCAGGAAGAGGCCAAGG  
 AGGAGGAACGCCTCATAGCTTTGAAGAAAATCCCTCAGCTCGGGGGTGTTCATTCTCATGGCCAAACGA  
 TGGGTCTCGCTTTGACCTGACTGTCCGCACGCAGAAGCAGGGGGATGACAGCTCTGAATGGGGGAACCTC  
 TTCTTCTGGAACCAGCTGTTGCACGTGCCCTTGAGGCAGCACCAGGTGAGCTGCTGTGACTGGCTGTGTA  
 AGATCATCTGCGGGGTGGTCACCATCCGCACCGTGTATGCCCTCCACAAGCAGGCCAAGGCCTGCCTCGT  
 CTCTCGCGTTAGCTGTGAGCGCACAGGCACCTCGCTTCCACACCCGTGGCGTGAACGACGACGGCCATGTG  
 TCCAACCTCGTGGAGACAGAGCAGATGATTTACATGGACGATGGAGTGTATCTTTTGTCCAGATCAGAG  
 GCTCCGTTCCGCTGTTCTGGGAACAGCCAGGGCTTCAGGTTGGCTCCCATCATCTGAGACTCCACAGAGG  
 CCTGGAAGCCAATGCCCCTGCTTTCGACAGGCACATGGTGTCTTGAAGGAGCAGTACGGGCAGCAGGTG  
 GTCGTGAACCTTCTGGGAAGCAGAGGCGGAGAGGAGGTGCTCAACAGAGCCTTCAAGAAGCTGCTCTGGG  
 CTTCTTGGCACGCGGGGCACACGCCCTATGATCAATTTTGACTTCCATCAGTTTGCCAAAGGTGGGAAGCT  
 AGAGAAATTTGAGAGCCCTCTTGAGGCCACAGTTAAAGCTGCACTGGGAAGACTTCGATGTGTTCAAAAG  
 GGGGAGAAGCTCAGTCCACGTTTTTCAGAAAGGCACCTTTCGGGATGAACGTCTTGACTGCCTGGACCGAA  
 CCAACACTGTGCAGAGCTTCATCGCGCTCGAGGTCTGCATCTGCAGCTCAAGACCCTGGGGCTGAGTTC  
 AAAACCCATCGTTGACCGCTTTGTGGAGTCTTCAAGCCATGTGGTCTCTGAATGGCCACAGCCTGAGC  
 AAGGTGTTACAGGCAGCAGAGCCCTGGAAGGGAAGGCCAAGGTGGGGAAGCTGAAGGATGGAGCCCGGT  
 CCATGTCTCGAACCATCCAGTCCAACCTTCTCGACGGGTGAAGCAGGAGGCCATCAAGCTGTCTGCTGGT  
 TGGGGACGTCTACGGCGAGGAGGTGGCAGACAAAGGGGGCATGCTGCTGGACAGCACGGCGCTCCTGGTG  
 ACTCCCAGGATCTGAAAGCTATGACTGAGCGTCAGTCCGAATTCACAAATTTCAAGCGGATCCGGATTG  
 CTATGGGGACCCTGGAACGTGAACGGAGGAAAGCAGTTCCGGAGCAACGTGCTCAGGACGGCGGAGCTGAC  
 AGACTGGCTGCTCGACTCGCCCCAGCTCTCGGGAGCTACCGACTCCCAGGATGACAGCAGCCCAGCTGAC  
 ATATTTGCTGTGGGGTTTGAAGAGATGGTGAATTGAGCGCAGGGAATATTGTCAATGCCAGTACTACCA  
 ACAAGAAGATGTGGGGTGAACAGCTTCAGAAAGCCATCTCACGCTCTCATAGATACATTCTGTTGACTTC  
 GGCACAGCTGGTGGGCGTCTGTCTTTATATCTTTGTACGTCCATACCATGTCCCGTTTCATCAGGGACGTA  
 GCCATCGACACAGTGAAGACGGGCATGGGGGGCAAGGCGGGGAACAAGGGCGCCGTCGGCATCCGCTTCC  
 AGTTCCACAGCACCAGCTTCTGCTTCATATGTAGTCACCTGACGGCCGGGCAGTCCCAGGTGAAGGAGCG  
 GAATGAAGACTACAAGGAGATCACCCAGAACTCTGCTTCCCAATGGGGAGAAATGTTTTTCTCATGAT  
 TATGTATTTTGGTGTGGCGATTTCAACTACCGCATTGATCTTACTTATGAAGAAGTCTTCTATTTTGTTA  
 AACGCCAAGACTGGAAGAACTTCTGGAATTTGATCAACTACAGCTACAGAAATCAAGTGGAAAAATTTT  
 TAAGGACTTTTACGAAGGAGCCATTAACCTTTGGACCCACCTACAAGTATGACGTTGGCTCAGCCGCCTAC  
 GATACAAGCGACAAATGCCGCACCCCGCTTGAGACAGCAGGGTGTGTGGTGGAGGAAGAACATCCCT  
 TTGATAAAACAGCTGGAGAACTCAACCTTCTAGACAGTGATCTAGATGTTGACACCAAAGTCAGACACAC  
 CTGGTCTCCTGGTGCCCTGCAGTATTATGGTCTGTGCGGAGCTACAAGCGTCTGATCACAGACCTGTGCTG  
 GCGATCGTGGAGGTGGAAGTTCAGGAAGTCGATGTGGGTGCTCGGGAGAGGGTTTTTCAGGAAGTGTCTT  
 CCTTCCAGGGCCCCCTGGATGCCACTGTTGTAGTAAACCTTCAATCACCGACCTTAGAAGAGAAAAACGA  
 GTTTCCAGAGGACCTGCGTACTGAGCTCATGCAGACCTTGGGGAGTTATGGGACAATTGTTCTTGTGAGG  
 ATCAACCAAGGGCAGATGCTGGTAACCTTTTGCAGACAGTCACTCGGCTCTCAGTGTCTTGGACGTGGACG  
 GTATGAAGGTGAAAGGCAGAGCAGTGAAGATTAGACCGAAGACCAAGGACTGGCTGAAAGGTTTGGGAGA  
 GGAGATCATTCGGAACGAGACAGCATGGCCCCGTGTCTCCCACTGCCAATCCTGTTTGTGCTGGAGGAA  
 AACTTTGACTTCACAAGTTTGGACTATGAGTCAGAAGGGGATATTCTTGAAGACGATGAAGACTACTTGG  
 TGGATGAATTCAATCAGCCTGGAGTCTCGGACAGTGAACCTCGGGGGAGACGACCTCTCTGATGTCCCCGG  
 CCCCACAGCACTGGCTCCTCCAGCAAGTCACCTGCTCTACCAAAAAGAAGCAGCATCCAACGTACAAA  
 GATGACGCGGACCTGGTGGAGCTCAAGCGGGAGCTGGAAGCCGTCGGGGAGTTCCGCCACCGTTCTCCGA  
 GCAGGTCTCTGTGGTCCCCAACCGGCTCGGCCACCTCAACCCCCGAGAGACCCCCCTCCAACCGG  
 TTTAATGGTGAAAAAGTCGGCTTCAGATGCGTCCATCTCCTCCGGCACCCATGGACAGTATTCAATTTTG  
 CAGACGGCAAGACTTCTACCAGGAGCACCTCAGCAACCTCCCAAGGCTCGGACTGGAATAAGTAAACCTT  
 ATAATGTCAAGCAGATCAAAACCACCAATGCCCAGGAGGCAGAAGCAGCAATCCGGTGTCTCCTGGAAGC  
 CAGAGGAGGTGCCTCCGAAGAAGCCCTAAGTGCCGTGGCCCCAAGGGACCTTGAAGCATCTCTGAACCA  
 GAGCCCACACGGGGGCAGCCAAACCAGAGACCCCAAGGCGCCCCACTCCTTCCCCGTGGGGCCCCAC  
 CCAGAGTTCTTGCATCAAGAAGCCAACCTTGAGAAGGACAGGAAGCCCTGTACACCGGAAGAAGAGTT  
 TGAGCAACAGACTGTCCATTTTACAATCGGGCCCCCGGAGACAAGCGTTGAGGCCCCCTCCTGTCTGTGACA  
 GCCCCCTCGAGTCCCTCTGTTCCCAAACCAAGAACATTTTCAGCCTGGGAAAGCTGCAGAGAGGCCAAGCC  
 ACAGGAAGCCAGCATCAGACGAAGCCCCCTCTGGGGCAGGAGCCTCTGTGCCACCACCTCTGGAGGCGCC  
 GCCTCTTGTGCCCAAGGTACCCCCGAGGAGGAAGAAGTCAGCCCCCGAGCCTTCCACCTGCAGGTCTCTG  
 CAGAGCAACAGCCAGCTTCTCCAGGGCCTCACTTACAATAGCAGTGACAGCCCCCTCTGGGCACCCACCTG  
 CCGCGGGCACCGTCTTCCACAAGGGGACTTTCTCAGCACTTCATCTGCTACAAGCCCCGACAGCGATGG  
 CACCAAAGCGATGAAGCCAGAGGCAGCCCCACTTCTTGGTGATTATCAGGACCCCTTCTGGAACCTTCTT

CACCACCC TAACTGTTGAATAACACTTGGCTTTCTAAGAGCTCAGACCC TTTGGACTCAGGAACCAGGA  
 GCCCCAAAAGAGATCCCATAGACCCAGTGTACAGCTGGCGCTTCAGCTGCCAAGGCAGAGCTGCCACCAGA  
 TCGTGAACACAAAACCTTAGGTCACTGGGTGACAATCAGTGACCAAGAAAAGAGGACAGCACTGCAGGTG  
 TTTGACCCACTGGCAAAAACATGACTGAGCAGCTTTGAAGGCTGCAGTCC TATAGAATGCATACCTTCCT  
 CCCTCTAGACATCCCTCCACCAGAAGAGACATCTATTTAAAGGCACACTGGCCAAAACGTTTGTGCATCT  
 GTCACCTCTCGTGTAGTTTACAAAAATCGTGTCTCTTATTCAGTAAGATGGTTACTCAGCCACCAAAATAT  
 ATTTCACTCAAGGCTTGTACATCTGAAGTTTGTCTTCAAGGAATGGGAACCTTCCTGTTAAATTCGGTG  
 TATGGATTTTAAAGAAAGGAATCTAGCCAATGAGGTCCAAGAAGTTCTCACCATTGAATTTTTAAATGGC  
 CATTTTGGGGAGCCACCTGAAGTTGATGTATAAAGTAAGGATTAGAGAAAAGAGGTCGTTGTGACCATT  
 ATAGCTGTCTCTGGCCCACTTAAACAAGGTTACAAAAAATCAGAGTCGGAAGCAGCCAAATAGGTCAACCT  
 AATGACTAGACTGTACATTTCCCATGAGCCTTCATGTTTAAAGTGTGTACATGTGCGTTAACCTTGATGATG  
 CGTGAATCCCGAGGGAGCCGGTGGCATAACCCGTTAGCTTAACTTAGCTTAACTAGCTGAAGGCTCCT  
 GTGCCATGCTTTAGACATTGCATGCCCTATCAATTACTATAATCC TGAGCCATGGTGTGCTACTGAAACC  
 AATTTTTATCCACCATCTAGTCTTTATTAATGAAACCTCACGGATCCTTTGTTCCGCTTATATTCCATG  
 CATACCACATAAAAGCACACAGTGCGAAAACCTTGCTGATACGCGATATTGATTCTCATTGTTAGAATA  
 TGGAGAGTGTTCAGCCTCGTCTGTCCGGCTGGAGCTTCGGGATGGAAAGTGCTATGTGTCCCTGCATAT  
 AAGAATCACCAGGCCAGTGTCTTGGGTTTGCTTGCTTATATGTTTGTCTATATTTTTTGCCTATACATT  
 TTTCCACGTTTCCAACAGCACTTCTCACCTATTCAATAACTGAAAAAGACATTACCATAGTGCTTTACA  
 TTTTTAAAGTAATGTTACAAGGCTTGAATCCATTTGGAGCAGATACCGTGTTCGCTATTTAATAAGA  
 AGTTCAGTAGTGAAATCTTACTGTACCGCTGTTGTATCTGGGAGCCTCGTACAGAGGCTCGCACAGCAG  
 TGATCAAGTGTCTATCCCTTACGTGACTGGGGGATGTCTGTCTTAAAGCTGACTGCTAGGATAGTAAGGA  
 TCATCTTGCCCTGGGCTATGCCACTGTCTTGTTACCAATTAGACATCTGGAATTTTATAATTAGTTTTTCAT  
 TGTCACCTGTCAAGATATATTGCAGATTACTTAAATATGGCCATCAAAACAAAAGTTACAACACGTATCTC  
 TTTTCATCTGAAAAC TAATACCTGGAAAAGGATAAAAAAAAAAAGGAATCCGTGACCCACAGAGCTAGAC  
 AGATAAGATGCATAGTTGACCAGTCATAAAAGGCGGTGTTTAAAGTGATCAGGATGCCGTTGGTGGCATT  
 ACGTGCTTTATATGATTTTTTACCTCTGTAACAAACACAAGAAATAAACAGAATGGTCTTTAACAGAGTTT  
 GGGGGAGAGAGCAAGATGGGTTCCTTGGAAGCTGATTTGCCAAGATGCACATCGCTATTAACAGCCAG  
 AGTCATAAATGAAATGAAATTGAAGAATTCATTCAAATGCTCTTTTCCCTATAACCTCTTTTCTCACCAA  
 AAAGGAGATAAATTTGAAAACAGATAAATGTAACAACAGTCAAAGAAGCAGGGGAAAAGTAAGCTCCTC  
 CAAAGTTGCTTGCAGTGTGGAATAGATCTATTTTATAGGTTTTCTCTTCGTTCCAGATACCAATAAA  
 TGGGACAGAGAATAAAAATTTTTGTTAAATATATGTGCTCATCTCCTAAGTAGCTCTTCAGAGTCTGACCGT  
 AAGTAAAAACACACAGAATTGTGTTGACTGGGGGAGGTGAATCACAAAAAGTTACGAGGAGTTTAAGAG  
 TTAAATATTATTTGATCGTGGCTGTCAAATTTAGTGAACAACATAGATTGGATTGGAGTTGGTAGTAAG  
 TATGGTTCTCATACCAGAATTCTCTT

>gi|37556102|ref|XM\_209367.2| Homo sapiens similar to dJ680N4.2  
 (ubiquitin-conjugating enzyme E2D 3 (homologous to yeast UBC4/5))  
 (LOC284767), mRNA

ATGGCGCTGAAACGGATTAATAAGGAACCTTAGTGATTTGGCCCGTGACCCCTCCAGCACAAATGTTCTGCAG  
 GTCCAGTTGGGGATGATATGTTTCACTGGCAAGCCACAATTATGGGACCTCATGACAGACCATATCAAGG  
 CGGTGTATTCTTTTGGACAATTCGTTTTCTACAGACTACCCCTTCAAACCATCTAAGGTTGCATTTACA  
 ACAAGAATTTATACCCCAAATATTAACAGTAATGGCAGCATTTTTCTTGACATTCTAAGATCACAGTGGT  
 CGCCTGCTTTAACACTTTCTAAAGTTCTTTTATCCATTGTTTCACTGCTATGTGATCCAAACCCAGATGA  
 TCCCTTAGTGCCAGAGATTGCACGGATCTATAAAACAGACAGAGATAAGTACAACAGAATATCTCGGGAA  
 TGGACTCAGAAGTATGCCATGTGA

>gi|19549332|ref|NM\_015983.2| Homo sapiens ubiquitin-conjugating enzyme  
 HBUCE1 (LOC51619), mRNA

GTGCAGTTGGTGGCGGTGAGCCGGCAGCGGGCCGCTCAGGCAGCCCCGGCCGGGCGCCCGGGTCCC  
 CGGCAGCGGGTAGGATGGCGCTAAAGCGGATCCAGAAGGAATTAACCGACTTGACAGAGGATCCTCCTG  
 CCCAGTGTCTGACAGACCTGTCTGGTGATGACTTGTTCCACTGGCAGGCCACCATCATGGGCCCCGAATGA  
 CAGTCTTTACCAAGGAGGTGTTTTCTTCTGACCATCCACTTTCTACAGATTACCGTTTCAAGCCCCCA  
 AAGGTTGCTTTACAACCAAAATTTATCACCTAATATCAACAGCAATGGCAGCATCTGCCTTGATATCC  
 TGCGGTCTCAGTGGTCTCCAGCGTTGACTGTGTCAAAGTTCTCTTGTCATCTGCTCGCTGCTCTGCGA  
 CCCCACCCCGATGACCCCTGGTGCCAGAGATAGCACACCTACAAGGCCGACAGAGAGAAGTACAAC  
 AGACTAGCAAGAGAGTGGACACAAAAATATGCTATGTAAGTGCTTGGAGGTTTTACATGAGACACTGTCT

CAAGAGAAGCTGGCAGAGAGGTCTTCCCTTAAAACCTTTGGGCTGTTGGCTGAGCCATTCAAAGAGCATCA  
TCTGTTCTTCAAACAAATGTTGGTCACCCACTCTCTCCAGCTGCAGCATGTTGGTGCCATTTTCAGCAAT  
TACGGCTTTGACAGTGCCACCTCTTTGATGCCAAATCAGCAACCATTGTTGTTATGATCTGCAGTCTTCC  
TGGTGACACTGGAATCTCTCTCTCTGCCGCCCTCAGTTTGTCTGCTGGTCTCTTGGGGGGCCAGGCCCTGC  
ACGTCTCTCCTACCCGGCCCTCAAATGGTGTCTTGCCTATGATGGTACCACACCAGGGCCTCAGCCTGGC  
CCCTCACCACATACCTTTTGCCTTTTAGAACTCAGTGCCATCCTGGGTGCCAGGGCAGAGCAGGCTTTG  
TTCGCACCTCATCTGCTGCAGAACACATCCTGAGGAGTCTCAGCTTATCCTGGAGGGAATTGGGAACAG  
TGTCAC TGGGAAGTGAAGGCC TAGCCCTGTGGCTTCCACCAGTCTCCTCCTGCAGTGCCACGTGGTGGCA  
TTTCTCGCCTCACACCAAGAAGCAGCAAGTGGAAAATTTTACAGGATACAAAGCACATAACACCCCATAGA  
GATGATTATGTTTTTAGAAGCAAGAGCAAAAATTTAGAACTCTAGAGATTTGGGTGATGTTACTCCATT  
TGATGAAGATTCTACTACCGCCCGCTCCTCCCATAGGAGCCTACACTAAGTCCAAGTGTGAGCCATTCA  
CAGACTAGAACACAAGGAGGGAGAGAGACTCTTAAACGTAAATAAAAATGCAATTCACTCACAACCTCT

>gi|33188457|ref|NM\_003339.2| Homo sapiens ubiquitin-conjugating enzyme  
E2D 2 (UBC4/5 homolog, yeast) (UBE2D2), transcript variant 1, mRNA  
GCTTCGCAGCGTCACGCCCTCCGGGGCCGTGGCGGCACGGCGGTGCGTAGCTTACTCACAGGGGGCGGCC  
CGTATCCCTCCGCCGCCGGCGCGGCTCGGCCCTCCCTCCCTGGCCCCGCAATCCCCGCGCCTCCCGACC  
TGCCCCCTCGGTGCGGGCCACCCCGTGTCTCCGACGGCCCCACCCGGCGGCGCAGCCCCGCCGCCGCCG  
TCCCTCGGTCCACCTGCAGCAGGGAGGAAGACAGGCAATCCCTCCGGCTGTCCGACCAAGAGAGGCCGGC  
CGAGCCCCGAGGCTTGGGCTTTTGTCTTCTGGCGGAGGGATCTGCGGCGGTTTAGGAGGCGGCGCTGATCC  
TGGGAGGAAGAGGCAGCTACGGCGGCGGCGGCGGTGGCGGCTAGGGCGGCGGCGAATAAAGGGGCCGCCG  
CCGGGTGATGCGGTGACCGCTGCGGCAGGCCAGGAGCTGAGTGGGCCCCGGCCCTCAGCCCGTCCCGCC  
GGACCCGCTTTCTCAACTCTCCATCTTCTCCTGCCGACCAGATCGCCGAGGCGGCCTCAGGCTCCCTA  
GCCCCTTCCCGTCCCTTTCCCCGCCCCCGTCCCCGCCCCGGGGGCCGCCGCCACCCGCCCTCCACCATGG  
CTCTGAAGAGAATCCACAAGGAATTGAATGATCTGGCACGGGACCTCCAGCACAGTGTTCAGCAGGTCC  
TGTTGGAGATGATATGTTCCATTGGCAAGCTACAATAATGGGGCCAAATGACAGTCCCTATCAGGGTGA  
GTATTTTCTTGACAATTCATTTCCCAACAGATTACCCCTTCAAACCACCTAAGGTTGCATTTACAACAA  
GAATTTATCATCCAAATATTAACAGTAATGGCAGCATTGTCTTTGATATTCTACGATCACAGTGGTCTCC  
AGCACTAACTATTTCAAAGTACTCTTGTCCATGTCTCTGTGTGTGATCCCAATCCAGATGATCCT  
TTAGTGCCGTAGATTGCTCGGATCTACAAAACAGATAGAGAAAAGTACAACAGAATAGCTCGGGAATGGA  
CTCAGAAGTATGCGATTAATTAAAGAAATTTAGGATAACCTCTACAAATAAAGATAGGGGAACCTGA  
AAGAGAAAGTCCTTTTGATTTCCATTTGACTGCTTTCTATGAGCCACGCCTCATCTTCCCCCTGTGCACA  
TGTTTACCTGATACAGCAGTGTGCGTGTGTGTGACATACCTGGAACAACAACTAGAAATACTGTACTTCT  
GTACCAACATTGCCTCCTAGCAGAGAAGTGTGTGTGTGACAAGCCAGTCTACAGGCATTACCTAGGTGT  
GAGACTAAAAGCTTTTCTTATTGACTTAAATTTGGATAACAGCAAGGTGTGAGGGGGGTGGTGGGTATGG  
TGTGTGCTTGGATGGGAAAGAAAAGGCTCCACTCACCTATAGGAGATTATTTTAAAGTGAATCCATTTA  
AACTCAAAACAGTTATGAAAAGCAAGGTGAAGAACATGAAGCTGTGTCTGTATTCTATTTTATTCCGAAGG  
AGCTACGCTTAGGTGAAAGTTATGACCAACCAGATTAACTCTACCCACATCCTGCATTTTAAAGGTCTA  
AGTTTAACTGGTCAACATTTAAATGGATTGGAGCTATTAGTACATCAAGTGTGATGGGCTTTGTTCCCAA  
CTCTTTTACATCTCCCTACCCCTTCAACCTTTGGCCTTTACGCCCTTCTTTCTCTTCCATATTCTTTG  
GTTTGTATGTGGTTTCTCAGTTAATACATAGCTAATAGCTCTTATTTTCTTATGTTTTTAAACCGCTTAG  
GTCTATTTGGATGTAAGGGTGAATTTCAATTTGATGGAATACCTTGTGTATATTTAAAGACCCAATTGCT  
CCTCTGGAGCTTGTACTTTCAAGAATGATTAATCTGTGTAATAAATGGTTACTACAGTCATTACATATA  
ATTTTGTGTGAATAGGCTTTTTTCATTTTAAAGATTTGTCTAGCTGAGATTAGTGGTGGATTTTCTCCC  
ACTTCTGAAATGTTTCAATTTTACTGGTTGCATTTTAAAGATCATGAAACAATTCAGTTACATTGTAAAAA  
GGATATCTTACGAGTAATTTTATTGAACAAGTTAGAGGCATAAGCTTAAAGAGCATTTCATGAAACAACA  
CATGCAGCATTCAGGAACCTTGATTGTTAAATTCATAAGAAATTTGCTTTATTAATGAAACTAAGCTGC  
ATTTTATCAAAACCTTGTGACATTCCCTTGGTACATAGGACATAAAACACAGAGGCATTGCTATTTGGTA  
AGTTAAGCTTCTGTGATTTGAATTTATAAAAGAGCAACATTGACCAAACTGGGAAACAAGAGCACAGCT  
TGTTTGGAGAGCTTACATAATTACTTTGCATTAACATTTGCAGGATGTTACACAATTTTAAATTTGACT  
GTATGTGGCTTTTGAAGTCTTCCCTTGACCCCTAGTAAAATATAGCTTGAAACTTGTAACAACCTGTGTT  
TGCCAGAAACATCATTCATGTGAAC TAGGCAAGTTACCTTTTTTCCCCCTTCTTTTCTAATTGTAAAC  
TAGGCCAACCTGAAAGCCATGGCTGATGCTCTAGCCATCAGGTCTTTCAAATGCATCTTTACACTCTTG  
CACAAAAGTTAAGGAATAAATGTCCACTGCTTTTGGTTTTTAAAAAAAAAAAAAAAAA

>gi|33149308|ref|NM\_003340.4| Homo sapiens ubiquitin-conjugating enzyme  
E2D 3 (UBC4/5 homolog, yeast) (UBE2D3), transcript variant 1, mRNA

ACCAAGTGAGGAACTGGGGGACGCTGTGGGGAGGGGCGTGGGGCTGGATCGCGCAGCGGCTGCTTCCTT  
 TACCTTCCCTCCCATGGTCTCCTTCCGGTTCTCGATGCTTCTCTGAGCCTAAGGGTTTCCGCCACTCGTTC  
 ACCCTCCCCCAGCTCATGATCTTCTCCTCCCTCCCCCGCCCTCCTGGTCCAATCTCCGATCTGTTTAGTAA  
 GAAGGTGCTGTTCCGAGAAGGAGAAGGAAAAGGGCTTGACACGTATTCACCTCGGCCCCGGACGTGGGAAG  
 CAAGCCGTCTGGCTTCGGCCTCACATCGGTCTGTGCTCGGGACGGCGGCGTGGCGGACTGATCCGCGG  
 CGGTGAAGAGGCGCCTGTGTCTGGCAGAGCTGGTGTGAGACGAGACAATCCTGCCCCGCCGCGGGATAA  
 TCAAGAGTTTGGCCGGACCTTTGAGCATACACCGAGAGAGTGAGGAGCCAGACGACAAGCACACACTAT  
 GGCCTGAAACGGATTAAATAAGGAACCTAGTGATTTGGCCCGTGACCTCCAGCACAAATGTTCTGCAGGT  
 CCAGTTGGGGATGATATGTTTCATTGGCAAGCCACAATTATGGGACCTAATGACAGCCCATATCAAGGCG  
 GTGTATTCTTTTGACAATTCAATTTCTTACAGACTACCCCTTCAAACCACCTAAGGTTGCATTTACAAC  
 AAGAATTTATCATCCAAATATTAACAGTAATGGCAGCATTTGTCTCGATATTCTAAGATCAGATGGTTCG  
 CCTGCTTTAACAATTTCTAAAGTTCTTTTATCCATTTGTTCACTGCTATGTGATCCAAACCAGATGACC  
 CCTAGTGCCAGAGATTGCACGGATCTATAAAACAGACAGAGATAAGTACAACAGAATATCTCGGGAATG  
 GACTCAGAAGTATGCCATGTGATGCTACCTTAAAGTCAGAATAACCTGCATTATAGCTGGAATAAACTTT  
 AAATTACTGTTCTTTTTTTGATTTTCTTATCCGGCTGCTCCCTATCAGACCTCATCTTTTTTAATTTTA  
 TTTTTTGTTTACCTCCCTCCATTCATTCACATGCTCATCTGAGAAGACTTAAGTTCTTCCAGCTTTGGAC  
 AATAACTGCTTTTAGAAACTGTAAAGTAGTTACAAGAGAACAGTTGCCCAAGACTCAGAATTTTTAAAAA  
 AAAAAATGGAGCATGTGTATTATGTGGCCAATGTCTTCACTCTAACTTGTTTATGAGACTAAAACCATTC  
 CTCACTGCTCTAACATGCTGAAGAAATCATCTGAGGGGGAGGGAGATGGATGCTCAGTTGTCACATCAAA  
 GGATACAGCATTATTTCTAGCAGCATCCATTCCTGTTTAAAGCTTCCACTGTTAGAGATTTGAGGTTACAT  
 GATATGCTTTATGCTCATAACTGATGTGGCTGGAGAATTGGTATTGAATTTATAGCATCAGCAGAACAGA  
 AAATGTGATGTATTTTATGCATGTCAATAAAGGAATGACCTGTTCTTGTCTACAGAGAATGGAAATTGG  
 AAGTCAAACACCCCTTTGTATTCCAAAATAGGGTCTCAAACATTTTGTAAATTTTCATTTAAATTGTTAGGA  
 GGCTTGGAGCTATTAGTTAATCTATCTTCCAATACACTGTTTAAATATAGCACTGAATAAATGATGCAAGT  
 TGTCATATGGATGAGTGATCAACTAATAGCTCTGCTAGTAATTGATTTATTTTTCTTCAATAAAGTTGCAT  
 AAACCAATGAGTTAGCTGCCCTGGATTAATCAGTATGGGAAACAATCTTTTGTAAATGCAAAGCTGTTTTT  
 TGTATATACTGTTGGGATTTGCTTCATTGTTTGACATCAAATGATGATGTAAAGTTGCAAAGAGTGAATA  
 TTTTGCCATGTTTCAAGTTAAAGTGCACAGTCTGTTACAGGTTGACACATTGCTTGACCTGATTTATGCAGA  
 ATTAATAAGCTATTTGGATAGTGTAGCTTTAATGTGCTGCACATGATACTGGCAGCCCTAGAGTTTCATAG  
 ATGGACTTTTGGGACCCAGCAGTTTGAATGTGTTTATGGAGTTTAAAGAAATTTATTTTCCAGGTGCAG  
 CCCCCTGCTTAACATGAAATTTCTCTTCACTTTGTACACTTGACAGCTGAAAAAAAACAACATGGGAGTAAT  
 AATGGGTCAAATTTGCAAAATAAAGTACTGTTTTGGTGTGGGAGTTGTCTATGAGGCTGTGTTGAAGTGA  
 CTTATCTATGTGGGATATTGAGTATCCATTGAAATGGATTTGTTTACGCCATTTACATTAATGAGCATTTA  
 AATGCAACAGATATCATTTTCAAGTGACTTAACATGAATGAATAAAAGTCAATGCTATTGGAAAAAAA  
 AAAAA

>gi|37539364|ref|XM\_353788.1| Homo sapiens similar to dopamine beta-hydroxylase-like (LOC378372), mRNA

ATGGCCCATGACCTTCTCTTCAGGCTTTTTCCACTTTTGGCCCTGGGAGTCCCCCTTACAAAGCAACCGCC  
 TTGGCCCCACATCTCGCCTGCGCTATTCCAGGTTCCCTAGATCCTTCTAATGTCAATTTTCTGCGTTGGGA  
 CTTTGACCTTGAGGCTGAAATCATCAGTTTTGAGCTCCAGGTCCGTACAGCTGGCTGGGTGGGCTTCGGT  
 GTCACAAATCGCTACACCAACGTGGGAAGTGATCTGGTTGTTGGAGGAGTCTTGCCTAATGGCAATGTCT  
 ATTTCTCGGATCAGCACTTGGTGGAGGAAGACACTCTGAAGGAGGATGGGAGCCAGGATGCTGAGCTGCT  
 GGGCTTGACGGAAGATGCTGTCTACACCACCATGCACTTTTCCAGGCCCTTCCGCTCCTGCGACCTCAT  
 GACCTGGATATTACGAGCAACACCGTGAGGGTGTGGCCGCCCTATGGCCTGGATGACACTCTGAAGCTGT  
 ATCGGGAGCGTACTTTTGTCAAGTCCATCTTTCTGTCTACAAGTCGTCCACCCTGACGATCTGGATGTCCC  
 TGAGGACACCATCATCCATGACTTGAGATCACTAATTTCTTCATTCCAGAGGATGACACCACGTATGCC  
 TGCACCTTTCTTCTCTCCCTATTGTGAGCGAGAAGCATCACATTTACAAGTTTGAGCCTAAGTTGGTCT  
 ACCACAATGAGACAACGGTGCATCACATCCTGGTGTATCGCCTGTGGCAATGCTAGCGTTCTCCCCACAGG  
 CATCAGCGACTGTATGGGGCCGACCTGCCCTTCTCCCTTTGCTCACAGGTATCGTGGGCTCGGCTGTC  
 GGGGGCACACTGTGCAAAATACGACACGGATGTCTCCAGCTGGGCTTCTTACGTTTTCCCATCCACTTCA  
 TCCCCCGGGCGCTGAGTCTTTCATGTCTTATGGGCTGTGTAGGACGGAGAAGTTTGAGGAGATGAATGG  
 AGCTCCTATGCCCTGACATACAGGTGTATGGCTACCTGCTGCACACCCACTTGGCTGGACGGGCTCTGCAA  
 GCAGTGCAATACAGGAATGGAACACAACCTTCAAAAAATCTGTAAAGACGATTCTTATGACTTCAACCTGC  
 AGGAGACTCGAGATTTGCCCTCTAGAGTGGAGATCAAGCCGGGAGATGAATTGCTGGTAGAATGTCTACTA  
 CCAGACACTGGACCGTGACTCCATGACATTTGGAGGCCCGAGCACCATTAAATGAGATGTGCTCATCTTC  
 CTCTTCTACTATCCCCAAAATAACATCTCCAGCTGCATGGGGTACCCTGACATTATCTACGTGGCCACG

AGCTAGGGGAGGAGGCATCAGAGTGA

>gi|24308084|ref|NM\_015529.1| Homo sapiens monooxygenase, DBH-like 1 (MOXD1), mRNA

TTTCCAAGGAAGGAAAGGCACACAATGGTATCAAATGTTTCATTCATTCATCTCTTGATGCTCTACGATA  
 TTATCAGTCTACACTATGCTTTTCTGAAAGGCCAGAAGTTCAAAGATGGACTAGTTTCCCAGGGACCTGA  
 TTCCCCAGTTGGAATACTCCAGCCCCTTGGAAATTCCCGGGATTATATAAAATAACTCTAGACAACAAGAC  
 TTTGTCTTTAAAGGTCCATGAATTCCTTTCTCTCTGTATTTAGGTATCCTGATTTTTCTTTTCCATATT  
 TTCCACAGGATTATTTTACAAATGCAAATAGAGAGTTGAAAAAGATGCTCAGCAAGATTACCATCTAGA  
 ATATGCCATGGAAAAATAGCACACACACAATAATTGAATTTACCAGAGAGCTGCATACATGTGACATAAAT  
 GACAAGAGTATAACGGATAGCACGTGTGAGAGTGATCTGGGCCCTACCACCATGAAGATGCAGGAGAAGCTG  
 GTCCCCAAGTACCATGACTCCAATAGGGGCACCAAGAGTTTGGCGTTATTGAATCCTGAGAAAAGTGTGT  
 GCTATCTACAGCCTTACCATACCTTGTATCTGGTAAATCAGGACGTCCCCATCCCAAACAAAGATACAACA  
 TATTGGTGCCAAATGTTTAAGATTCTGTGTTCGAAGAAAGCATCATGTAATAAAGGTTGAGCCAGTGA  
 TACAGAGAGGCCATGAGAGTCTGGTGCACCACATCTGCTCTATCAGTGCAGCAACAACCTTAAACGACAG  
 CGTTCTGGAGTCCGGCCACGAGTGCTATCACCCAACATGCCCGATGCATTCTCACCTGTGAAACTGTG  
 ATTTTTGCCCTGGGCTATTGGTGGAGAGGGCTTTTCTTATCCACCTCATGTTGGATTATCCCTTGGCACTC  
 CATTAGATCCGCATTATGTGCTCCTAGAAGTCCATTATGATAATCCCACTTATGAGGAAGGCTTAATAGA  
 TAATTCCTGGACTGAGGTTATTTTACACAATGGATATAAGGAAATATGATGCTGGGGTGATTGAGGCTGGC  
 CTCTGGGTGAGCCTCTTCCATACCATCCCTCCAGGGATGCCCTGAGTTCCAGTCTGAGGGTCACTGCACCTT  
 TGGAGTGCCCTGGAAGAGGCTCTGGAAGCCGAAAAGCCAAGTGGAATTCATGTGTTTGCTGTTCTTCTCCA  
 TGCTCACCTGGCTGGCAGAGGCATCAGGCTGCGTCAATTTTGAAAAGGGAAGGAAATGAAATTACTTGCC  
 TATGATGATGATTTTGACTTCAATTTCCAGGAGTTTCAGTATCTAAAGGAAGAACAACAATCTTACCAG  
 GAGATAACCTAATTACTGAGTGTCGCTACAACACGAAAGATAGAGCTGAGATGACTTGGGGAGGACTAAG  
 CACCAGGAGTGAAATGTGTCTCTCATACCTTCTTTATTTACCCAAGAATTAATCTTACTCGATGTGCAAGT  
 ATTCAGACATTATGGAACAACCTCAGTTTCATTGGGGTTAAGGAGATCTACAGACCAGTCACGACCTGGC  
 CTTTCATTATCAAAAGTCTCAAGCAATATAAAAACCTTTCTTTTCATGGATGCTATGAATAAGTTTAAATG  
 GACTAAAAAGGAAGGTCCTCTTCAACAAGCTGGTCCCTCAGCTGCCAGTGAATGTGAGATGTTTCAAG  
 ACAGACAATGCTGAGTGGTCGATTCAAGGAATGACAGCATTACCTCCAGATATAGAAAGACCTATAAAG  
 CAGAACCCTTTGGTGTGTGGCAGCTTCTTCTCTTCCCTGCACAGAGATTTCTCCATCAACTGCTTGT  
 TTGCCCTTCTGCTACTGACGTGCACGCTGAGCACCAAGAGCTTGTGATCAAAATTCTGTTGGACTTGACAA  
 TGTTTTCTATGATCTGAACCTGTCTATTTGAAGTACAGGTTAAAGACTGTGTCCACTTTGGGCATGAAGAG  
 TGTGGAGACTTTTCTTCCCCATTTTCCCTCCCTCCCTTTTCTCTTCCATGTTACATGAGAGACATCAATC  
 AGGTTCTCTTCTCTTTCTTAGAAATATCTGATGTTATATATACATGGTCAATAAAATAAACTGGCCTGA  
 CTTAAGATAACCATTTTAAAAAATTTGGGCTGTCTGTTGGGAATAAAAGAATTCTTTCTTTCTTACTACAT  
 TCTGTTTTATTTAAATACTCATTTGTTGCTATTTTCACTTTTGTGACTTTTATATTTCTTTAAAAAAT  
 TCCTTCTTTTAAAAAATATAAAAGGGACTACTGTTTCATTCCAGTTTTCTTCTTCTTTGTTGTTCTTCTA  
 GTGTGACTTTTCAAGTGTAACAGCCATTCTTCTTGACTTTAATATTGTCCAGTTCTGGTCTTTTCTGTGA  
 ATTACCACGGGCCCCCTTACCCTAATGCTTTTTGTTGATGCCCACTCTGGTTCCCTTGTATTCTGAGTC  
 TGTTGGTACCCCAAATGACCCACACCCATYTTAAAGTACTTTTTTTTACCTTCCCTGTTTGTAGTACTGGC  
 CAGATGAGTTTTTTCTAGAGCTCTGTCTACTATCTGAAAAGAAAGAGGCTATGGGAAACATAGAAATGGTA  
 TGTATTAATAACTGATCATAGGCTGAGGAGAAAAATGTAGCTGGCTGCAAACCCAGTGCTGTGAGGTGA  
 CTTATATGAGGTTCCAGATCAAAGACAGGCCGTGTGAGCCAGTCCAGGAGGGTGTAAAGTTCTGAATGGTT  
 CCTTGCTGACTTTGGGTGACACATGTACCACATACTGGCTCAGTTTAAAGTCATGGTTCTATTGTAGATTT  
 ATTTTTATATTAGTTAATAAATGACTTTAAATTGTACCAATTGAAAATCTTGTCACTCTTTTGGTTTTCT  
 TTTATATAGCTCAGCCAAATCTCTGTTTTATGTCTGTCTCATCTCTTAAGCTAAATCTGTTTGGATCA  
 TATTAATAAACCTCGTGCCG

>gi|40254422|ref|NP\_000594.2| nitric oxide synthase 3 (endothelial cell) [Homo sapiens]

MGNLKSVAQEPGPPCGLGLGLGLGLCGKQGPATPAPEPSRAPASLLPPAPEHSPSSPLTQPPEGPKFPR  
 VKNWEVGSITYDTLSAQQAQDGPCTPRRCLGSLVFPRLQGRPSGPPAPEQLLSQARDFINQYYSSIKR  
 SGSQAHEQRLQEVEAEVAATGTYQLRESELVFGAKQAWRNAPRCVGRIQWGLQVFDARDCRSAQEMFTY  
 ICNHIKYATNRGNLRSATVFPQRCPRGRDFRIWNSQLVRYAGYRQQDGSVRGDPANVEITELCIQHWGT  
 PGNGRFDVLPLLLQAPDDPPELFLLPPELVLEVPLEHPTLEWFAALGLRWYALPAVSNMLLEIGGLEFPA

APFSGWYMSTEIGTRNLCDPHRYNILEDVAVCMDDLTRTTSSLWKDKAAVEINAVLHSHYQLAKVTIVDH  
 HAATASFMKHLNEQKARGGC PADWAWIVPPI SGLTPVFHQEMVNYFLSPA FRYQDPDPWKGSAAKGTGI  
 TRKKTTFKEVANAVKI SASLMGTVMKRVKATILYGSETGRAQSYAQQLGRLFRKAFDPRVLCMDEYDVVS  
 LEHETLVLVVTSTFGNGDPPENGESFAAALMEMSGPYNSSPRPEQHKSXYKIRFNSISCS DPLVSSWRRKR  
 KESSNTDSAGALGTLRFVFLGSRAYPHFCAFARA VDTREELGGERLLQLGQGDELGCQEEAFRGWAQ  
 AAFQAACETFCVGEDAKAAARDIFS PKRSWKQRQYRLSAQAEGQLLPGLIHVHRRKMFQATIRSVENLQ  
 SSKSTRATILVRLDTGGQEGLOYQPGDHIGVCP PNRPGLV EALLSRVEDPPAPTEPVAVEQLEKGSPPGP  
 PPGWVRDPRLPCTLRQALTFFLDITSPSPQLLRLSTLAE EPREQEQEALSDQPRRYE EWKWFRCPT  
 LLEVLEQFPSPVALPAPLLLTLQLPLLQPRYYSVSSAPSTHPGEIHLTVAVLAYRTQDGLGPLHYGVCSTWL  
 SQLKPGDPVPCFIRGAPSFRLPPDPSLPC ILVGP GTGIAPFRGFQERLHDIESKGLQPTPMTLVFGCRC  
 SQLDHLRDEVQNAQQRGVFGRVLTAFSREP DNPKTYVQDILRTELAAEVHRLVCLERGHMFVCGDVTMA  
 TNVLQTVQRILATEGDMELDEAGDVIGVLRDQQR YHEDIFGLTLRTQEVTSRIRTQSFSLQERQLRGAVP  
 WAFDPGSDTNSP

>gi|4506061|ref|NP\_002724.1| protein kinase, AMP-activated, gamma 1  
 non-catalytic subunit; AMPK gamma 1; Protein kinase, AMP-activated,  
 noncatalytic, gamma-1 [Homo sapiens]  
 METVISSDSSPAVENEHPQETPESNNSVYTSFMKSHRCYDLIPTSSKL VVFDTS LQVKKAFFALVTNGVR  
 AAPLWDSKKQSFVGMLTITDFINILHRYYKSALVQIYELEEHKIETWREVYLQDSFKPLVCISP NASLFD  
 AVSSLIRNKIHRLPVIDPESGNTLYILTHKRILKFLKFITEFPKPEFMSKSLEELQIGTYANIAMVRTT  
 TPVYVALGIFVQHRVSALPVVDEKGRVVDIYSKFDVINLAAEKTNNLDVSVTKALQHRSHYFEGVLKCY  
 LHETLETIINRLVEAEVHRLVVVDENDVVKGIVSLSDILQALVLTGGEKKP

>gi|33186925|ref|NP\_057287.2| protein kinase, AMP-activated, gamma 2  
 non-catalytic subunit; H91620p protein [Homo sapiens]  
 MGSAMVMDTKKKKDVSSPGSGGKKNASQKRRSLRVHIPDLSSFAMPLLDGDLEGSGKHSSRKVDSPFGPG  
 SPSKGGFSRGPQPRPSSPMSAPVRPKTSPGSPKTVFPFSYQESPPRSPRMSFSGIFRSSSKESSPN SNP  
 ATSPGGIRFFSRSRKTSGLSSSPSTPTQVTKQHTFPLESYKHEPERLENRIYASSSPDGTGQRF CSPSFQ  
 SPTRPPLASPTHYAPSKAAALAAALGPAEAGMLEKLEFEDEAVEDSESGVYMRFMRS HKCYDIVPTSSKL  
 VVFDTTLQVKKAFFALVANGVRAAPLWESKKQSFVGMLTITDFINILHRYYKSPMVQIYELEEHKIETWR  
 ELYLQETFKPLVNI SPDASLFDVYSLIKNIHRLPVIDPISGNALYILTHKRILKFLQFLMSDMPKPAF  
 MKQNLDELIGITYHNI AFIHPTPI IKALNIFVERRISALPVVDESGKVVDIYSKFDVINLAAEKTNNL  
 DITVTQALQHRSQYFEGVVKCNKLEILETIVDRIVRAEVHRLVVVNEADSIVGII SLSDILQALILTPAG  
 AKQKETETE

>gi|8394044|ref|NP\_059127.1| protein kinase, AMP-activated, gamma 3  
 non-catalytic subunit [Homo sapiens]  
 MEPGLEHALRRTPSWSSLGGSEHQEMS FLEQENSSSWPSPAVTSSSERIRGKRRAKALRWTRQKSVEEGE  
 PPGQGEGRSRPAAESTGLEATFPKTTPLAQADPAGVGTPPTGWDCLPSDCTASAAGSSTDDVELATEFP  
 ATEAWECELEGLLEERPALCLSPQAPFPKLGWDEL RKPGAQIYMRFIEHTCYDAMATSSKLVI FDTML  
 EIKKAFFALVANGVRAAPLWDSKKQSFVGMLTITDFILVLHRYYSPLVQIYEIEQH KIETWREIYLQGC  
 FKPLVSI SPNDSLFEAVYTLIKNRIHRLPVLDPVSGNVLHILTHKRLLKFLHIFGSLLP RPSFLYRTIQD  
 LGIGTFRDLAVVLETAPILTALDIFVDRRV SALPVVNECGQVVGLYSRFDVIHLAAQQTYNHLDMSVGEA  
 LRKRTLCLGVLSCQPHESLGEVIDRIAREQVHRLVLVDETQHLLGVVSLSDILQALVLS PAGIDPSGPE  
 KI

>gi|24041029|ref|NP\_000616.3| nitric oxide synthase 2A isoform 1; NOS,  
 type II; nitric oxide synthase, macrophage [Homo sapiens]  
 MACPWKFLFKTKFHQYAMNGEKDINNVEKAPCATSSPVTQDDLQYHNLSKQQNESPQPLVETGKKSPES  
 LVKLDATPLSSPRHVRIKNWGSGMTFQDTLHHKAGGILTCRSKSLGSI MTPKSLTRGPRDKPTPPDELL  
 PQAIEFVNQYYGSFKEAKIEEHLARVEAVTKEIETTGT YQLTGDELIFATKQAWRNAPRCIGRIQWSNLQ  
 VFDARSCSTAREMF EHCIRHVRYSTNNGNIRS AITVFPQRS DKGKDFRVWNAQLIRYAGYQMPDG SIRGD  
 PANVEFTQLCIDLGWKPKYGRFDVVPVLVLQANGRDPELFEI PPDLVLEVAMEHPKYEFWRELELKWYALP  
 AVANMLLEVGGLFPGCPFN GWYMGTEIGVRDFCDVQRYNILEEVGRMGLETHKLASLWKDQAVVEINI  
 AVLHSFQKQNVTIMDHSSAAESFMKYMQNEYRSRGGC PADWIWLVPPMSG SITPVFHQEMLN YVLSPFYY  
 YQVEAWKTHVWQDEKRRPKRREIPLKVLVKAVL FACMLMRKTMASRV RVTILFATETGKSEALAWDLGAL  
 FSCAFNPKVVCMDKYRLSCLEERLLLVTSTFGNGDCPGNGEKLKKS LFMKLNNKFRYAVFGLGSSM



YPRFCAFAHDIDQKLSHLGASQLTPMGEGDELSGQEDAFRSWAVQTFKAACETFDVRGKQHIQIPKLYTS  
 NVTWDPHHYRLVQDSQPLDLKALSSMHAKNVFTMRLKSRQNLQSPTSSRATILVELSCEDGQGLNYLPG  
 EHLGVC PGNQPALVQGILERVVDGPTPHQTVRLEALDESGSYWVSDKRLPPCSLSQALTYFLDITTPPTQ  
 LLLQKLAQVATEEPERQRLALCQPSEYSKWKFTNSPTFLEVLEEFPSLRVSAGFLLS QLPILKPRFYSI  
 SSSRDHTPTIEHILTAVVVTYHTRDGQGPLHHGVCSTWLNLSLKPQDPVPCFVRNASGFHLPEDPSHPCILI  
 GPGTGIAFFRSFWQQLHDSQHKGVVGRMTLVFGCRPDEDHIYQEEMLEMAQKGVLAHVHTAYSRLPG  
 KPKVYVQDILRQQLASEVLRVLHKEPGHLYVCGDVRMARDVAHTLKQLVAAKLKLNEEQVEDYFFQLKSQ  
 KRYHEDIFGAVFPYEAKKDRVAVQPSLSLEMSAL

>gi|4504789|ref|NP\_002211.1| 1D-myo-inositol-trisphosphate 3-kinase A  
 [Homo sapiens]

MTLPGGPTGMARPGGARPCSPGLERAPRRSVGELRLLFEARCAAVAAAAAAGEPRARGAKRRGGQVPNGL  
 PRAPPAPVPIQLTVTAEEPDPVPTSPGPPERERDCLPAAGSSHLQPPRLSTSSVSSTGSSSLLEDSEDD  
 LLSDSESRSRGNVQLEAGEDVQKNHWQKIRTMVNLPVISPFKKRYAWVQLAGHTGSFKAAGTSGILILKR  
 CSEPERYCLARLMADALRGCVPAFHGVVERDGE SYLQLQDLLDGFDPGVLDCKMGVVRTYLEEELTKARE  
 RPKLRKDMYKMLAVDPEAPTEEEHAQRAVTKPRYMQRWREGISSSTTLGFRIEGIKKADGSCSTDFKTR  
 SREQVLRVFEFVQGDDEVLRRYLNRLQQIRDTELEVSEFFRRHEVIGSSLLFVHDHCHRAGVWLIDFGKT  
 TPLPDGQILDHRRPWEEGNREDGYLLGLDNLIGILASLAER

>gi|38569400|ref|NP\_002212.2| 1D-myo-inositol-trisphosphate 3-kinase B;  
 IP3 3-kinase [Homo sapiens]

MAVYCYALNSLVIMNSANEMKSGGGPGPSGSETPPPPRAVLSPGVSFSPGRGASFLFPPAESLSPEEPR  
 SPGGWRSGRRRLNSSSGSGSGSSSSVSSPSWAGRLRGDRQQVVAAGTLPSPGPEEAKRKLRLQRELQN  
 VQVNQKVGMEFAHIIQAQSSAIQAPRSRLGRARSPSPC PFRSSSQPPGRVLVQGARSEERRTKSWGEQCP  
 ETSGTDSGRKGGPSLCSQVKKGMPLPGRAAPTGSEAQGSAFVRMEKGIPASPRCGSPTAMEIDKRG  
 PTPGTRSLAPSLGLFGASLTMAEVAARVTSTGPHRPQDLALTEPSGRARELEDLQPPEALVERQGGFL  
 GSETSPAPERGGPRDGEPPGKMGKGYLPCGMPGSGEPEVGKRPEETTVSVQSAESSDSLWSRLPRALAS  
 VGPEEARS GAPVGGGRWQLSDRVEGGSPTLGLLGGSPSAQPGTGNVEAGIPSGRMLEPLPCWDAADLKE  
 PQCPGGRVGVQPGNSRVWQGTMEKAGLAWTRGTGVQSEG TWESQRQSDALPSELLPDQDQKPFRLKA  
 CSPSNIPAVIITDMGTQEDGALEETQGS PRGNLPLRLKLSSSASSTGFSSSYEDSEDISSDPERTLDPN  
 SAFLHTLDQKPRVSKSRKIKNMVHWSPFVMSFKKYPWIIQLAGHAGSFKAAANGRIKHKHCESEQRCL  
 DRLMVDVLRPFVPAYHGDVVKDGERYNQMDLLADFDSPCVMCKMGIRTYLEELTKARKKPSLRKDMY  
 QKMIEVDPEAPTEEEKAQRAVTKPRYMQRWRETISSTATLGFRIEGIKKEDGTVNRDFKKTREQVTEAF  
 REFTKGNHNILAIYRDLKAIRTTELVSPFFKCHEVIGSSLLFIHDKKEQAKVWMIDFGKTTPLPEGQTL  
 QHDVPWQEGNREDGYLSGLNNLVDILTEMSQDAPLA

>gi|18643383|ref|NP\_079470.1| inositol 1,4,5-trisphosphate 3-kinase C  
 [Homo sapiens]

MRRPCRCGSLNEAEAGALPAAARMGLEAPRGRRRQPGQQRPGPGAGAPAGRPEGGGPWARTEGSSSLHSE  
 PERAGLGPA PGTESPQAEFWTDGQTEPAAAGLVETERPKQKTEPDRSSLRTHLEWSWSELETTCLWTET  
 GTDGLWTDPHRSDLQFQPEEAS PWTQPGVHG PWTELETHGSQTQPERVKSWADNLWTHQNSSSLQTHPEG  
 ACPSKEPSADGSWKELYTDGSRTQQDIEGPWTEPYTDGSQKKQDTEAARKQPGTGGFQIQDQTDGSWTQP  
 STDGSQTAPGTDCLLGEPEDEGPLEEPEPGELLTHLYSHLKCSPLCPVPRLIITPETPEPEAQPVGPPSRV  
 EGGSGGFSSASSFDESEDDVAGGGGASDPEDRSGSKPWKKLKTVLKYSPFVVSFRKHYPWVQLSGHAGN  
 FQAGEDGRILKRFCQCEQRSLEQLMKDPLRPFPVAYYGMVLQDQGT FNQMEDLLADFEGPSIMDCKMGS  
 TYLEELVKARERPRPRKDMYEKMAVDPGAPTPEEHAQGAVTKPRYMQRWRETMSSTSTLGFRIEGIKKA  
 DGTCTNTNFKKTQALEQVTKVLEDFVDGDHVLQKYVACLEELREALEISPFFKTHEVVGSSLLFVHDHTG  
 LAKVWMIDFGKTVLALPDHQTLSHRLPWAEGNREDGYLWGLDNMICLLQGLAQS

>gi|4758698|ref|NP\_002363.1| mannosidase, alpha, class 2A, member 1;  
 mannosidase, alpha type II [Homo sapiens]

MKLSRQFTVFGSAIFCVVIFSLYLMLDRGHLDPNPRREGSFPQGQLSMLQEKIDHLERLLAENNEIIS  
 NIRDVINLSESVEDGPKSSQSNFSQAGSHLLPSQLSLSDTADCLFASQSGSHNSDVQMLDVYSLISF  
 DNPDPGGVWKQGF DITYESNEWDTEPLQVFVPHSHNDPGWLKTFNDYFRDKTQYIFNNMVLKLKEDSRRK  
 FIWSEISYLSKWWDIIDIQKKDAVKSLIENGQLEIVTGGWMPDEATPHYFALIDQLIEGHQWLENNIGV  
 KPRSGWAIDPFGHSPTMAYLLNRAGLSHMLIQRVHYAVKKHFALHKTLEFFWRQNWDLGSVTDILCHMMP  
 FYSYDIPHTCGPDPKICCFDFKRLPGGRFGCPWGVPETIHPGNVQSRARMLLDQYRKKSKLFRTKVLL



APLGDDFRYCEYTEWDLQFKNYQQFLDYMNSSQSKFKVKIQFGTSLDFFDALDKAETQRDKGQSMFPVLS  
 GDFFTYADRDDHYWSGYFTSRPFYKRMDRIMESHLRAAEILYYFALRQAHKYKINKFLSSSLYTALTEAR  
 RNLGLFQHHDAITGTAKDWVVDYGTRLFHSLMVLEKIIIGNSAFLILKDKLTYDSYSPDTFLEMDLKQK  
 SQDSLPLQKNIIIRLSAEPYLVVYNPLEQDRISLVSVYVSSPTVQVFSASGKPEVQVSAVWDTANTISET  
 AYEISFRAHIPPLGLKVYKILESASSNSHLADYVLYKNKVEDSGIFTIKNMINTEEGITLENSFVLLRFD  
 QTGLMKQMMTKEDGKHHEVNVQFSWYGTTIKRDKSGAYLFLPDGNAKPYVYTTPPFVRVTHGRIYSEVTC  
 FFDHVTHRVRLYHIQIGIEGQSVEVSNIVDIRKVYNREIAMKISSDIKSQNRFYTDLNGYQIQPRMTLSKL  
 PLQANVYPMTTMAYIQDAKHRLTLLSAQSLGVSSLSNGQIEVIMDRRLMQDDNRGLEQGIQDNKITANLF  
 RILLEKRSVANTEEEEKSVSYPSLLSHITSSLMNHPVIPMANKFSPTLELQGEFSPLQSSSLPCDIHLVNL  
 RTIQSKVGNGHSNEAALILHRKGFDCRFSSKGTGLFCSTTQGKILVQKLLNKFIVESLTPSSLSLMHSPP  
 GTQNISEINLSPMEISTFRIQLR

>gi|3123244|sp|P49641|M2A2\_HUMAN Alpha-mannosidase IIx (Mannosyl-  
 oligosaccharide 1,3-1,6-alpha-mannosidase) (MAN IIx) (Mannosidase alpha  
 class 2A member 2)

MKLKKQVTVCGAAIFCVAVFSLYLMLDRVQHDPTRHQNGGNFPRSQISVLQNRIEQLEQLLEENHEIISH  
 IKDSVLELTANAEGPPAMLPHYTVNGSWVVPPEPRPSFFSISPDQCQFALGGRGQKPELQMLTVSEELPF  
 DNVDGGVWRQGFDISYDPHDWAEDLQVFVPHSHNDPGWIKTFDKYYTEQTQHILNSMVSKLQEDPRRR  
 FLWAEVSFFAKWWDNINVQKRAAVRRLVGNGQLEIATGGWMPDEANSHYFALIDQLIEGHQWLERNLGA  
 TPRSGWAVDPFGYSSTMPYLLRRANLTSMLIQRVHYAIKKHFAATHSLEFMWRQTWSDSDSTDFCHMMP  
 FYSYDVPHTCGPDPKICCFDFKRLPGGRINCPWKVPRAITEANVAERAALLLDQYRKKSQLFRSNVLL  
 VPLGDDFRYDKPQEWDAQFFNYQRLFDFFNSRPNLHVQAQFGTSLDYFDALYKRTGVEPGARPPGFPVLS  
 GDFFSYADREDHYWTGYYSRPFYKSLDRVLEAHLRGAEVLYSLAAAHARRSGLAGRYPLSDFTLLTEAR  
 RTLGLFQHHDAITGTAKAVVDYGVRLRLSLVNLKQVIIHAAHYLVLGDKETYHFDPEAPFLQVDDTRL  
 SHDALPERTVIQLDSSPRFVVLNPLEQERFSMVSLLVNSPRVRVLSSEEQPLAVQISAHWSSATEAVPD  
 VYQVSVVRLPALGLGLVQLQLGLDGHRTLPSSVRIYLHGRQLSVSRHEAFPLRVIDSGTSDFALSRYM  
 QVWFSGLTGLLKSIRRVDEEHEQVDMQVLVYGTRTSKDKSGAYLFLPDGEASPTSPRSPCCVSLKALS  
 SQRWLRMTSTFTRRSFGFTICQGWGCLWTYHPWWTSGTTSTRSWPCTSIQTSTARVQPRRYLKKLPLOAN  
 FYPMFVMAYIQDAQKRLTLHTAALGVSSSLKDGQLEVIDRRLMQDDNRGLGQGLKDNKRTCNRFRLLE  
 RRTVGSEVQDSHSTSYPSLLSHLTSMYLNAPALPVARMQLPGPGLRSFHLPLASSLPCDFHLLNLRTLQ  
 AEEDTLPSAETALILHRKGFDCGLEAKNLGFNCTTSQGKVALGSLFHGLDVVFLQPTSLTLTYPLASPSN  
 STDVYLEPMEIATFRLRLG

>gi|10835173|ref|NP\_000611.1| nitric oxide synthase 1 (neuronal) [Homo  
 sapiens]

MEDHMFVGQQIQPNVISVRLFKRKVGGLGFLVKERVSKPPVVISDLIRGGAAEQSGLIQAGDIILAVNGR  
 PLVDLSYDSALEVLRGIASETHVVLILRGPEGFTTHLETTFTGDGTPKTIRVTQPLGPPTKAVDLSHQPP  
 AGKEQPLAVDGASGPGNGPQHAYDDGQEAGSLPHANGLAPRPPGQDPAKKATRVSLQGRGENNELLKEIE  
 PVLSSLTSGSRGVKGGAPAKAEMKDMGIQVDRDLGKSHKPLPLGVENDRVFNDLWGKGNVPVVLNNPYS  
 EKEQPPTSGKQSPTKNGSPSKCPRFLKVKNWETEVVLTDTLHLKSTLETGCTEYICMGSIMHPSQHARRP  
 EDVRTKGQLFPLAKEFIDQYYSSIKRFGSKAHERLEEVENKEIDTTSTYQLKDTIELIYGAKHAWRNASRC  
 VGRIQWSKLQVFDARDCTTAHGMFNYICNHVKYATNKGNLRSAITIFPQRTDGKHDFRVWNSQLIRYAGY  
 KQPDGSTLGDPANVQFTEICIQQGWKPPRGRFDVLPPLLQANGNDPELFQIPPELVLEVPPIRHPKFEWFK  
 DLGLKWYGLPAVSNNMELLEIGGLEFSACPFSGWYMGTEIGVRDYCDNSRYNILEEVAKKMNLDMRKTSSLW  
 KDQALVEINIAVLYSFQSDKVTIVDHHSATESFIKHMEYRCRGCPADWVWIVPPMSGSIPTVFHQEM  
 LNYRLTPSFEYQPDPNWTHVWKGTTNGTPTKRRRAIGFKKLAEAVKFSAKLMGQAMAKRVKATILYATETGK  
 SQAYAKTLCEIFKHAFDAKVMSEEDIVHLEHETLVLVVTSTFGNGDPPENGEKFGCALMEMRHPNSVQ  
 EERKSYKVRFNSSVSSYSDSQSSGDGPDRLDNFESAGPLANVRFSVFGLSRAYPHFCAFGHAVDTLLEE  
 LGGERILKMREGDELCEGQEEAFRTWAKKVFKAACDVFVCGDDVNIEKANNSLISNDRSWKRNKFRLTFFVA  
 EAPLETQGLSNVHKRVSAARLLSRQNLQSPKSSRTIFVRLHTNGSQELQYQPGDHLGVFPGNHEDLVN  
 ALIERLEDAPPVNMVVKVELLEERTALGVISNWTDELRLPPCTIFQAFKYLDITTPPTPLQLQQFASL  
 ATSEKEKQRLVLVLSKGLQEYEEWKWGKNPTIVEVLEEFPSIQMPATLLLTQLSLLQPRYYSISSSPDMYP  
 DEVHLTVAIVSYRTRDGEPIHGVCSWLNRIQADELVPCFVRGAPSFHLPRNPQVPCILVPGGTGIAP  
 FRSFQQRRQFDIQHKGMNCPMVLVFGCRQSKIDHIYREETLQAKNKGVFRELYTAYSREPKPKKYVQD  
 ILQEQLAESVYRALKEQGGHIYVCGDVTMAADVLAQIRIMTQQGKLSAEDAGVFI SRMRDDNRYHEDIF  
 GVTLRTYEVTNRLRSESI AFIEESKKTDEVFSS

>gi|37540252|ref|XP\_291315.2| similar to KIAA1815 protein [Homo sapiens]

MEWGESAAVRRHRVGVVERREGAAAAPPEREARAQEPLVDGCSGGGRTRKRSPPGGSGGASRGAGTGLSE  
VRAALGLALYLIALRTLVLQSLQQLVLRGAAGHRGEFDALQARDYLEHITSIGPRTTGSPENEILTIVHYL  
LEQIKLIEVQSNLSHKISVDVQRPTGSFSIDFLGGFTSYDNTNIVVKLEPRDGAQHAVLANCHFDSVA  
NSPGASDDAVSCSVMLEVLRLVSTSSSEALHHAIVFLFNGAEENVLQASHGFIQHPWASLIRAFINLEAA  
GVGGKELVFQTGPENPWLQAYVSAAKHPFASVVAQEVFQSGIIPSDTDFRIYRDFGNIPGIDLAFIENG  
YIYHTKYDTADRILTDSIQRAGDNILAVLKHLATSDMLAAASKYRHGNMVFDFVLGLFVIAYPRIIGSII  
NYMVMGVVLYLGKKFLQPKHKTGNKKDFLCGLGITLISWFTSLVTVLIIAVFISLIGQSLSWYNHFYV  
SVCLYGTATVAKIILIHITLAKRFYYMNASAQYLGEVFFDISLFVHCCFLVTLTYQGLCSAFISAVWVAF  
LLTKLCVHKDFKQHGAKGKFIAYFLLGMFIPYLYALYLIWAVFEMFTPIGRSGSEIPDPVVLASILAGC  
TMLSSYFINFIYLAKESTKKTMLTLTLVCAITFLLVCSGTFPPYSSNPANPKPKRVFLQHMTRTFHDLEG  
NAVKRDSGIWINGFDYTGISHITPHIPEINDSIRAHCEENAPLCGFPWYLPVHFLIRKNWYLPAPVSPR  
NPPHFRLLISKEQTPWDSIKLTFEATGPSHMSFYVRAHKGSTLSQWSLGNPTVTSKGGDYFVFYSHGLQA  
SAWQFWIEVQVSEEHPEGMVTVAIAAHYLSGEDKRSPLDALKEKFPDWTFPSAWVCTYDLFVF

>gi|21361382|ref|NP\_009102.2| protein-O-mannosyltransferase 1 [Homo sapiens]

MWGFLKRPVVVTADINLSLVALTGMGLLSRLWRLTYPRVVVFDEVYQGQYISFYMKQIFFLDDSGPPFGH  
MVLALGGYLGFGDGNFLWNRIGAEYSSNPVWSLRLPALAGALSVPMAYQIVLELHFSHCAAMGAALLM  
LIKNALITQSRMLLESVLIFNLLAVLSYLKFFNCQKHSFSLSWFWLTLTGACSCAVGIKYMVGFT  
YVLVLGVAHVHAWHLLGDQTLNVLGADVQCCMRPACMGQMRMSQGVCFCHLLARAVALLVIPVLYLLF  
FYVHLILVFRSGPHDQIMSSAFQASLEGGLARITQGGPLEVAFGSQVTLRNVFGKPVPCWLHSHQDTPM  
IYENGRGSSHQQVTCYPFKDVNNWVIVKDPRRHQLVSSPPRVRHGMVQLVHGMTTRSLNTHDVAAP  
LSPHSQEVSCYIDYNISMPAQNLRLEIVNRGSDTDVWKTILSEVRFVHVNTSAVLKLSGAHLPDWGYRQ  
LEIVGEKLSRGYHGSTVWNVEEHRYGASQEQRERERELHSPAQVDVSRNLSFMAFSELOQRMLALRSD  
SEHKYSSSPLEWVTLDTNIAWHLHPRTSAQIHLLGNIVIVWSGLALAIYALLSLWYLLRRRRNVHDL  
DAWLRLWLAGALCAGGWAVNYLPFFLMEKTLFLYHYLPALTFQILLPVVLQHISDHLCRSLOQRNSFSA  
LVVAWYSSACHVSNTRLPLTYGDKSLSPHELKALRWKDSWDILIRKH

>gi|4506321|ref|NP\_002837.1| protein tyrosine phosphatase, receptor type, N precursor; islet cell antigen 2; islet cell antigen 512; islet cell autoantigen 3; protein tyrosine phosphatase-like N precursor [Homo sapiens]

MRRPRRPGGLGGSGGLRLLCLLLSSRPGGCSAVSAHGCLFDRRLCSHLEVCIQDGLFGQCQVGVGQAR  
PLLQVTSPLVQLRQLQGLRQLMSQGLSWHDDLQYVISQEMERIPRLRPPEPRPRDRSGLAPKRPAGEL  
LLQDIPTGSAPAAQHRLPQPPVGKGGAGASSLSPLQAEELLPLLEHLLPPQPPHPSLSYEPALLQPYL  
FHQFGSRDGSRVSESGPMVSVGPLPKAEAPALFSRTASKGIFGDHPGHSYGDLPGPSAPQLFQDSGLLY  
LAQELPAPSRARVPRLEQGGSSRAEDSPEGYEKEGLGRGEKPASPAVQPDALQRLAAVLAGYGVELR  
QLTPEQLSTLLTLLQLLPGKAGRNPGGVVNVGADIKKTMEGPVEGRDTAELPARTSPMPGHPTASPTSSE  
VQQVPSVPSSEPPKAARPPVTPVLEKKSPLGQSQPTVAGQPSARPAEEYGYIVTDQKPLSLAAGVKLL  
EILAEHVHMSSGSFINISVVGPAITFRIRHNEQNLSLADVTQQAGLVKSELEAQTGLQILQTGVGQREEA  
AAVLPPQTAHSTSPMRSVLLTLVALAGVAGLLVALAVALCVRQHARQQDKERLAALGPEGAGDPTTFEYQD  
LCRQHMAKSLFNRAEGPPEPSRVSSVSSQFSDAAQASPSHSSSTPSWCEEPAQANMDISTGHMILAYME  
DHLNRNDRDLAKEWQALCAYQAEPTNCATAQEGEKNRHPDFLPYDHARIKLKVESSPSRSDYINASPI  
IEHDPRMPAYIATQGPLSHTIADFWMVWESGCTVIVMLTPLVEDGVKQCDRYWPDEGASLYHVYEVNLV  
SEHIWCEDFLVRSFYLNKVTQETRTLQFHFLSWPAEGTPASTRPLLDFFRRKVNKCYRGRSCPIIVHCS  
DGAGRTGTYYILDMVLNRMAKGVKEIDIAATLEHVRDQRPGLVRSKDQFEFALTAVAEVNAAILKALPQ

>gi|11386149|ref|NP\_002838.1| protein tyrosine phosphatase, receptor type, N polypeptide 2 isoform 1 precursor; protein tyrosine phosphatase receptor pi; phogrin; tyrosine phosphatase IA-2 beta; IAR/receptor-like protein-tyrosine phosphatase [Homo sapiens]

MGPPLPLLLLLLLLLLPPRVLPAPSSVPRGRQLPGRLGCLLEGLCGASEACVNDGVFGRQCQKVPAMDFY  
RYEVSVALQRLRVALQKLSGTGFTWQDDYTQYVMDQELADLPKTYLRRPEASSPARPSKHSVGSERRYS  
REGGAALANALRRHLPFLEALSQAPASDVLARTHTAQDRPPAEGDDRFSESILTYVAHTSALTYPGPRT  
QLREDLLPRTLGLQLPDELSPKVDSGVDRHLLMAALSAYAAQRPPAPPGEGLSEPQYLLRAPSRMPRPLL

APAAPQKWPSPLGDSSEDPSTGDGARIHTLLKDLQRQPAEVRGLSGLELDGMAELMAGLMQGVHDGVARG  
 SPGRAALGESGEQADGPKATLRGDSFPDDGVQDDDDRLYQEVHRLSATLGGLLQDHGSRLLPALPFARP  
 LDMERKKSEHPSSLSSEEETAGVENVKSQTYSKDLLGQQPHSEPGAAAFGELQNQMPGPSKEEQSLPAG  
 AQEALSDGLQLEVQPSEEEARGYIVTDRDPLRPEEGRRLLVEDVARLLQVPSSAFADVEVLGPAVTFKVSA  
 NVQNVTTEDVEKATVDNKKLEETSGLKILQTVGSGSKSLKFLPPQAEQEDSTKFIALTLVSLACILGVL  
 LASGLIYCLRHSSQHRLKEKLSGLGGDPGADATAAYQELCRQRMATRPDRPEGPHTSRISVSSQFSDG  
 PIPSPSARSSASSWSEEPVQSNMDISTGHMILSYMEDHLKNKNRLEKEWEALCAYQAEPNSSFVAQREEN  
 VPKNRSLAVLTYDHSRVLLKAENSHSHSDYINASPIMDHDPNRPAYIATQGPLPATVADFWQMVWESGCV  
 VIVMLTPLAENGVRQCYHYWPDGNSLYHIYEVNLVSEHIWCEDFLVRSFYLNQTNETRTVTQFHFLS  
 WYDRGVPSSSRSLDFRRKVNKCVRGRSCPIIVHCSDGAGRSQTYVLIDMVLNKMAGKAKEIDIAATLEH  
 LRDQRPGMVQTKEQFEFALTAVAEVNAAILKALPQ

>gi|4507335|ref|NP\_003886.1| synaptojanin 1; inositol 5'-phosphatase  
 (synaptojanin 1); synaptojanin-1, polyphosphoinositide phosphatase  
 [Homo sapiens]

MAFSKGFRIYHKLDPFSLIVETRHKKECLMFESGAVAVLSSAEKEAIKGTYSKVLDAYGLLGVLRLNL  
 GDTMLHYLVLTGCMVSGKIQESEVFRVTSTEFISLRIDSSDEDRISEVRKVLNSGNFYFAWSASGISLD  
 LSLNAHRSMQEQTDDNRFFWNQSLHLHLKHYGVNCDWLLRLMCGGVEIRTIYAAHKQAKACLISRLSCE  
 RAGTRFNVRGTTNDGHVANFVETEQQVYLLDSDVSSFIQIRGSVPLFWEQPGQVGSVHRVMSRGFEANAP  
 AFDRHFRTLKNLYGRQIIVNLLGSKEGEHMLSKAFQSHLKASEHAADIQMVNFYHQMVKGGKAELHVS  
 LKPQVQKFLDYGYFFYNGSEVQRCQSGTVRTNCLDCLDRNTSVQAFGLGLEMLAKQLEALGLAEKPQLVTR  
 FQEVFRSMWSVNGDSISKIYAGTGALGKAKLDGARSVTRTIQNNFFDSSKQEAIDVLLGLNTLNSDLA  
 DKARALLTTGSLRVSEQTLQSASSKVLKSMCENFYKYSKPKKIRVCVGTWNVNGGKQFRSIAFKNQTLTD  
 WLLDAPKLAGIQEFQDKRSKPTDIFAIGFEEMVELNAGNIVSASTTNQKLWAVELQKTI SRDNKYVLLAS  
 EQLVGVCFLVFIRPQHAPFIRDVAVDVTVKTMGGATGNKGAVAIRMLFHTTSLCFVCSHFAAGQSQVKER  
 NEDFIEIARKLSFPMGRMLFSDYVFWCGDFNYRIDLPNEEVKELIRQONWDSLIAGDQLINQKNAGQVF  
 RGFLEGKVTFAPTYKYDLFSDDYDTSEKCRTPAWTDRVLWRRRKWPFDRSAEDLDLLNASFQDESKILYT  
 WTPGTLHYGRAELKTS DHRPVVALIDIDIFEVEAEERQNIYKEVIAVQGPDPGTVLVSIKSSLPENNFF  
 DDALIDELLQQFASFGEVILIRFVEDKMWVTFLEGSSALNVLSLNGKELLNRTITIALKSPDWIKNLNEEE  
 MSLEKISIALPSTSTSLTGEDA EVAADFMEGDVDDYSAEVEELLQPQLPSSSSGLTSPSSSPRTSP  
 CQSPTISEGFPVPSLPIRPSRAPSRTPGPPSAQSSPIDAQPATPLPKDPAQPLEPKRPPPPRPVAPPTRP  
 APPQRPSPPSGARSPAPTRKEFGGIGAPSPGVARREMEAPKSPGTTTRKDNIGRSQSPQAGLAGPGPAG  
 YSTARPTIPPRAGVISAPQSHARASAGRLTPESQSKTSETSKGSTFLPEPLKPQAAFPQSSLPPPAQRRL  
 QEPLVPVAAMPQSGPQPNLETPPQPPPSRSSHSLPSEASSQPQVKTNGISDGKRESPLKIDPFEDLSF  
 NLLAVSKAQLSVQTSVPVTPDPKRLIQLPSATQSNVNTLSSVSCMPTMPPIPARSQSQENMRSSPNPFIT  
 GLTRTNPFSDRTAAPGNPFRAKSESEATSWFSKEEPTVISPFPQLPLGHNKSRASSSLDGFKDSFDLQ  
 GQSTLKISNPKGWVTFEEEEDFGVKGKSKSACSDLLGNQPSFSGSNLTLNDDWNKGTNVSFVCLPSRRP  
 PPPPVPLLPPTSPVPDPFTTLASKASPTLDFTER

>gi|26190608|ref|NP\_003889.1| synaptojanin 2; inositol phosphate 5'-  
 phosphatase 2 (synaptojanin 2) [Homo sapiens]

MALSKGLRLGLRLGAEGDCSVLLEARGRDCLLFEAGTVATLAPEEKEVIKQYQKLTDAYGCLGELRLK  
 SGGTSLSFLVLVTGCTSVGRIPDAEIIKITATDFYPLQEEAKEEERLIALKKILSSGVFYFSWPNDGSRF  
 DLTVRTQKQGDSSSEWGNSSFFWNQLLHVPLRQHQSVCDDWLLKIIICGVVTIRTVYASHKQAKACLVSRVS  
 CERTGTRFHTRGVNDDGHVSNFVETEQMIYMDGVSFVQIRGSVPLFWEQPGQVGSVHRLRLHRLGLEAN  
 APAFDRHMLVLLKEQYQGVVNNLLGSRGEEVLNRAFKLLWASCHAGDTPMINFDFHQFAKGGKLEKLE  
 TLLRPQLKLHWEDFDVFTKGENVSPRFQKGTLMNCLDCLDRNTVQSFIALEVLHLQLKTLGLSSKPIV  
 DRFVESFKAMWSLNGHSLSKVFTGSRALEGKAKVGNLKDGAARMSTIQSNFFDGVKQEAIKLLLVGDVY  
 GEEVADKSGGMILLDSTALLVTPRILKAMTERQSEFTNFKRIRIAMGTWNVNGGKQFRSNVLRITAEITDWLL  
 DSPQLSGATDSQDDSPADIFAVGFEEVELSAGNIVNASTTNKMWGEQLQKAI SRSHRYILLTSAQLV  
 GVCLYIFVRPHYVPFIRDVAIDTVKTMGGKAGNKGAVGIRFQFHSTSFICSHLTAGQSQVKERNEDY  
 KEITQKLCFPMGRNVFSDYVFWCGDFNYRIDLTYYEEVFYVVKRQDWKKLLEFDQLQLQKSSGKIFKDFH  
 EGAINFGPTYKYDVGSAAYDTSKCRTPAWTDRVLWRRKKHPFDKTAGELNLLSDLDVDTKVRHTWSPG  
 ALQYYGRAELQASDHRPVLAIVEVEVQEVQVGVARERVFQEVSSFGQPLDATVVVNLQSPITLEEKNEFPED  
 LRTELMQTLGSYGTIVLVRINQGQMLVTFADSHSALSVLDVDMKVKGRAVKIRPKTKDWLKGLEEIIR  
 KRDSMAPVSPTANSCLLEENFDFTSLDYSEGDILEDDDEDYLVDEFNQPGVSDSELGGDDLSDVPGPTAL  
 APPSKSPALTKKKQHPTYKDDADLVELKRELEAVGEFRHRSPSRSLSVPNRPRPPQPPQRPPTGLMVK

KSASDASISSGTHGQYSILQTARLLPGAPQQPPKARTGISKPYNVKQIKTTNAQEAEAAIRCLLEARGGA  
 SEEALSAVAPRDLEASSEPEPTPGAAPKPTPQAPPLLPRRPPPRVPAIKKPTLRRTGKPLSPREEQFEQQT  
 VHFTIGPPETSVEAPPVVTAPRVPPVPKPRTFQPGKAAERPSHRKPASDEAPPGAGASVPPPLEAPPLVP  
 KVPRRRKKSAPAAFHLQVLQSNSQLLQGLTYNSSSDSPSGHPPAAGTVFPQGDFLSTSSATSPDSDGTKAM  
 KPEAAPLLGDYQDPFWNLLHHPKLLNNTWLSKSSDPLDSGTRSPKRPIDPVSAGASAAKAELPPDHEHK  
 TLGHWVTISDQEKRTALQVFDPLAKT

>gi|27485630|ref|XP\_209367.1| similar to dJ680N4.2 (ubiquitin-  
 conjugating enzyme E2D 3 (homologous to yeast UBC4/5)) [Homo sapiens]  
 MALKRINKELSDLARDPPAQCSAGPVGDDMFHWQATIMGPHDRPYQGGVFFWTIRFPTDYPFKPSKVAFT  
 TRIYHPNINSNGSIFLDILRSQWSPALTLKSVLLSICSLLCDPNPDDPLVPEIARIYKTDKYNRISRE  
 WTQKYAM

>gi|8393719|ref|NP\_057067.1| ubiquitin-conjugating enzyme HBUCE1 [Homo  
 sapiens]  
 MALKRIQKELTDLQRDPPAQCSAGPVGDDLFHWQATIMGPNDSPYQGGVFFLTIFHPTDYPFKPPKVAFT  
 TKIYHPNINSNGSICLDILRSQWSPALTVSKVLLSICSLLCDPNPDDPLVPEIAHTYKADREKYNRLARE  
 WTQKYAM

>gi|33188456|ref|NP\_862821.1| ubiquitin-conjugating enzyme E2D 2  
 isoform 2; ubiquitin carrier protein; ubiquitin-conjugating enzyme E2-  
 17 kDa 2 [Homo sapiens]  
 MFHWQATIMGPNDSPYQGGVFFLTIFHPTDYPFKPPKVAFTTRIYHPNINSNGSICLDILRSQWSPALTI  
 SKVLLSICSLLCDPNPDDPLVPEIARIYKTDREKYNRIAREWTQKYAM

>gi|33149322|ref|NP\_871621.1| ubiquitin-conjugating enzyme E2D 3  
 isoform 2; ubiquitin-conjugating enzyme E2-17 kDa 3; ubiquitin carrier  
 protein [Homo sapiens]  
 MALKRINKELSDLARDPPAQCSAGPVGDDMFHWQATIMGPNDSPYQGGVFFLTIFHPTDYPFKPPKVAFT  
 TRIYHPNINSNGSICLDILRSQWSPALTISKVLLSICSLLCDPNPDDPLVPEIARIYKTDKYNRLARE  
 WTEKYAML

>gi|37539365|ref|XP\_353789.1| similar to dopamine beta-hydroxylase-like  
 [Homo sapiens]  
 MAHDLLFRLFPLLALGVPLQSNRLGPTSRRLYSRFLDPSNVIFLRWDFDLEAEIISFELQVRTAGVWGFG  
 VTNRYTNVGSDDLVGGLVPLNGNVYFSDQHLVEEDTLKEDGSQDAELLGLTEDAVYTTMHFSRPFRCSDPH  
 DLDITSNTVRVLAAYGLDDTLKLYRERTFVKISIFLLQVHPDDLDVPEDTIIHDLEITNFLIPEDDTTYA  
 CTFLPLPIVSEKHIIYKFEPKLVYHNETTVHHILVYACGNASVLPTGISDCYGADPAFSLCSQVIVGSAV  
 GGTLCKYDSDLVLQLGFFTFPIHFIPGAESFMSYGLCRTEKFEEMNGAPMPDIQVYGYLLHHLAAGRALQ  
 AVQYRNGTQLRKICKDSDYDFNLQETRDLP SRVEIKPGDELLVECHYQTLDRDSMTFGGPSTINEMCLIF  
 LFYYPQNNISSCMGYPDIIYVAHELGEASE

>gi|37183305|gb|AAQ89452.1| dopamine-oxygenase [Homo sapiens]  
 MCCWPLLLLWGLLPGTAAGGSGRTYPHRTLLDSEGKYWLGSQRGSQIAFRLQVRTAGYVGFSGFSPTGAM  
 ASADIVVGGVAHGRPYLQDYFTNANRELKKDAQDYHLEYAMENSTHTIIIEFTRELHTCDINDKSITDST  
 VRVIWAYHHEDAGEAGPKYHDSNRGTSKSLRLNPEKTSVLSTALPYFDLVNQDVPIPNKDTTYWCQMFKI  
 PVFQEKHHVIKVEPVIQRGHESLVHHILLYQCSNNFNDSVLES GHECYHPNMPDAFLT CETVIFAWAIGG  
 EGFSYPHPVGLSLGTPLDPHYVLLLEVHYDNPTYEEGLIDNSGLRFLYTM DIRKYDAGVIEAGLWVSLFHT  
 IPPGMPEFQSEGHCTLECLEEAEAEKPSGIHVFAVLLHAHLAAGRIRLRHFRKGEMKLLAYDDDFDFN  
 FQEFQYLKEEQITILPGDNLITECRYNTKDRAEMTWGGLSTRSEMCLSYLLYYPRINLRCAISIPDIMEQL  
 QFIGVKEIYRPVTTWPFIIKSPKQYKNLSFMDAMNKFKWTKEGLSFNKLVLVSLPVNVRCSKTDNAEWSI  
 QGMTALPPDIERPYKAEPLVCGTSSSSSLHRDFSINLLVCLLLLSTLSTKSL